

# Late Triassic (Tuvalian – Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones) ostracods from Monte Gambanera (Castel di Iudica, Central-Eastern Sicily, Italy)

Sylvie Crasquin<sup>1,\*</sup>, Francesco Sciuto<sup>2,\*</sup>, Agatino Reitano<sup>3</sup> and Rosa Maria Coco<sup>2</sup>

<sup>1</sup> CR2P – CNRS, MNHN, Sorbonne Université, 4, Place Jussieu, T. 46-56, E.5, F-75252 Paris Cedex 05, France

<sup>2</sup> Palaeoecological Research Group, Department of Biological, Geological and Environmental Science, Catania University, Corso Italia, 55, Catania 95129, Italy

<sup>3</sup> Natural History Museum, via degli Studi 9, 97013 Comiso, Ragusa, Italy

Received: 28 October 2019 / Accepted: 2 July 2020 / Published online: 22 December 2020

**Abstract** – Ostracod associations coming from the Upper Triassic (*Tropites subbullatus*/*Anatropites spinosus* zones of the Tuvalian substage) clays and sandstones of the Mufara Formation outcropping along the west side of Monte Gambanera (Castel di Iudica, central-eastern Sicily) have been analysed for the first time. The specimens are relatively abundant, silicified, well preserved and often preserved as complete carapaces. Over 200 specimens have been determined. They belong to the families Healdiidae, Bairdiidae, Bythocyprididae, Acrafiidae, Cytheruridae, Limnocytheridae, Candonidae, Cavellinidae, Polycopidae and Thaumatoocyprididae. Thirty-seven species are identified of which nine species are new: *Hungarella forelae* n.sp., *Hungarella siciliensis* n.sp., *Bairdia andreocrasquini* n.sp., *Bairdia gambaneraensis* n.sp., *Ptychobairdia iudicaensis* n.sp., *Ptychobairdia leonardo* n.sp., *Petasobairdia jeandercourti* n.sp., *Kerocythere dittainoensis* n.sp. and *Mockella barbroae* n.sp.

**Keywords:** Ostracods / systematic / Late Triassic / Neo-Tethys / Central-Eastern Sicily / Mount Gambanera / Mufara Formation

**Résumé** – Ostracodes du Trias supérieur (Tuvalien, Carnien, zones à *Tropites subbullatus*/*Anatropites spinosus*) de Monte Gambanera (Castel di Iudica, Sicile Centre Est, Italie). Pour la première fois est ici analysée une association d'ostracodes provenant du Trias supérieur (zones à *Tropites subbullatus*/*Anatropites spinosus* du sous étage Tuvalien) dans les argiles et grès de la Formation Mufara affleurant le flanc ouest du Mont Gambanera (Castel di Iudica, Sicile Centre Est). Les spécimens, silicifiés, sont relativement abondants, bien préservés et les plus souvent retrouvés sous formes de carapaces complètes. Plus de 200 spécimens ont été identifiés. Ils appartiennent aux familles Healdiidae, Bairdiidae, Bythocyprididae, Acrafiidae, Cytheruridae, Limnocytheridae, Candonidae, Cavellinidae, Polycopidae and Thaumatoocyprididae. Trente-sept espèces sont reconnues dont sont nouvelles: *Hungarella forelae* n.sp., *Hungarella siciliensis* n.sp., *Bairdia andreocrasquini* n.sp., *Bairdia gambaneraensis* n.sp., *Ptychobairdia iudicaensis* n.sp., *Ptychobairdia leonardo* n.sp., *Petasobairdia jeandercourti* n.sp., *Kerocythere dittainoensis* n.sp. and *Mockella barbroae* n.sp.

**Mots clés :** ostracodes / systématique / Trias supérieur / Néo-Téthys / Sicile centre est / Mont Gambanera / Formation Mufara

## 1 Introduction


This is the second contribution on the Late Triassic ostracod fauna of the Mufara Formation outcropping in central eastern Sicily. Previously the ostracod fauna of the *Tropites dilleri*

zone of the Tuvalian substage outcropping at Monte Scalpello has been analysed (Crasquin *et al.*, 2018).

Now, a sedimentary level which is stratigraphically higher than the previous one and referable to the *Tropites subbullatus*/*Anatropites spinosus* zones of the Tuvalian substage, has been identified at the western side of the Monte Gambanera, nine kilometres south of Monte Scalpello (Fig. 1). The samples provided a rich and mostly well-preserved ostracod fauna.

\*Corresponding author:

[sylvie.crasquin@sorbonne-universite.fr](mailto:sylvie.crasquin@sorbonne-universite.fr)




**Fig. 1.** Geographical location of Monte Gambanera, Sicily, Italy and sample locality.

## 2 Geological setting and studied samples

Monte Gambanera is a modest relief located in central eastern Sicily (F 269 III NE of the Carta d'Italia alla scala 1:25 000) to the southeast of the town of Castel di Iudica (EN), about 40 kilometres west of Catania (Fig. 1). Structurally Monte Gambanera is part of the “Monte Judica Units” (Lentini *et al.*, 1987) and is inserted along the northern margin of the Gela Foredeep, in the geodynamic context of the southern end of the Maghrebian–Sicilian Southern Apennine nappes (Lentini *et al.*, 1987; Grasso, 2001 *inter alia*s).

In the Monte Gambanera area, the outcropping sediments are assigned to the Neo-Tethyan Mesozoic–Cenozoic complex which belongs to the so-called Imerese Succession (Lentini *et al.*, 1987; Montanari, 1987; *inter alia*s) or Imerese-Sicano Succession (Carrillat and Martini, 2009; Di Paolo *et al.*, 2012). The Imerese Basin, where these sedimentary successions were deposited, was delimited by the Panormide Carbonate Platform to the west and the Trapanese Carbonate Platform to the east and south (Catalano and D’Argenio, 1982; Montanari, 1987; Speranza and Minelli, 2014). In this basin, therefore, occurs a transitional facies between the Panormide and Trapanese shelf facies, on the one hand, and a deep marine facies of the Neo-Tethys, on the other. The sedimentary succession of Monte Gambanera (Fig. 2) starts with the “Carnian Flysch” (Auct.) or the Mufara Formation (Schmidt di Friedberg and Trovò, 1962). This unit, outcropping in the



**Fig. 2.** Stratigraphic series of Monte Gambanera, Sicily, Italy.

southern slopes of the mount, mainly consists of dark grey pelites, which locally contain rare ammonites, with rare interbeds of fossiliferous calcarenites and fibrous calcite with *Halobia* spp. imprints. The Mufara Fm. has been analysed since the beginning of the nineteenth century by Calcara (1840, 1845), Nelli (1899a, b) and subsequently by Gemmellaro (1904), Scalia (1907a, b, 1909, 1910–1914), Maugeri Patanè (1934) and Lentini (1974). These latter authors attributed these sediments to the Carnian (Late Triassic).


Stratigraphically, the Mufara Fm. outcropping at Monte Scalpello, can be referred to the *Tropites dilleri* zone of the Tuvalian substage (Crasquin *et al.*, 2018) due to the presence of Trachyceratidae (*Neoprotrachyceras*, *Trachysagenites*, *Pamphagosirenites*) and Tropitidae. A foraminifera, conodont and palynomorph biostratigraphical analysis, allowed to attribute the levels of the Mufara Fm. outcropping at Monte Gambanera to the *Tropites subbullatus/Anatropites spinosus* zones of the Tuvalian substage (Fig. 3) (Carrillat, 2001; Carrillat and Martini, 2009). In this stratigraphic horizon, cropping out near masseria Balconere at the west side of Mount Gambanera, two levels consisting of slightly silty clays have been sampled (Fig. 1). As they are stratigraphically very close and the number of specimens is quite low, we consider here the ostracod assemblages of both samples in all. They represent the ostracod association of the present study.

Twenty kilograms of sediments were collected from each of the two stratigraphic levels. Sediments were routinely washed, dried in oven and sieved. Then, ostracod specimens were picked out from the 63  $\mu\text{m}$  fraction. The ostracod specimens were examined and measured under a stereomicroscope, then photographed under an LMU Tescan Vega II SEM. The material is housed in the Palaeontological Museum of the University of Catania. The repository number of the specimens are given in the systematic descriptions and/or in plate explanations. Over 200 specimens have been picked out from the two samples. The specimens are silicified, quite well preserved and often consist of complete carapaces.

## 3 Systematic palaeontology

**Abbreviations.** L: length; H: height; W: width; RV: right valve; LV: left valve; DB: dorsal border; VB: ventral border; AB: anterior border; PB: posterior border; PVB: postero-ventral border; AVB: antero-ventral border; PDB: postero-





**Plate 1.** Ostracods from Late Triassic (Tuvalian-Carnian) of Monte Gambanera, Sicily, Italy. All the specimens are stored in the Palaeontological Museum of the University of Catania. The repository numbers are given as PCM (Palaeontological Museum Catania) O (Ostracods) X H (number of holotype) or X P (number of paratype) or FS X (Figured Specimen number) registration date. Scale bars = 200  $\mu$ m. A: *Hungarella forelae* n.sp. A: holotype, right lateral view of a complete carapace, PMC O 21 H13/10/2019; paratype figured in Figure 6A (Crasquin *et al.*, 2018). B-C: *Hungarella siciliensis* n.sp. B: holotype, right lateral view of a complete carapace, PMC O 22 H 13/10/2019; C: paratype, right lateral view of a complete carapace, PMC O 78 P 13/10/2019. D: *Hungarella* sp. A: right lateral view of a complete carapace, PCM O FS49. E-F: *Bairdia andreocrasquini* n.sp. E: holotype right lateral view of a complete carapace, PMC O 23 H 13/10/2019; F: paratype, right lateral view of a complete carapace, PMC O 79 P 13/10/2019. G-H: *Bairdia gambaneraensis* n.sp. G: holotype, right lateral view of a complete carapace, PMC O 24 H 13/10/2019; H: paratype, right lateral view of a complete carapace, PMC O 80 P 13/10/2019. I: *Bairdia cassiana* (Reuss, 1869). Right lateral view of a complete carapace, PCM O FS50. J: *Bairdia* cf. *monostorii* Forel and Grădinaru (2018). Right lateral view of a complete carapace, PCM O FS51. K: *Bairdia* sp. 1 *sensu* Crasquin *et al.* (2018). Right lateral view of a complete carapace, PCM O FS52. L: *Bairdia* sp. A. Right lateral view of a complete carapace, PCM O FS53. M: *Bairdia* sp. B. Right lateral view of a complete carapace, PCM O FS54. N: *Bairdia* sp. C. Right lateral view of a complete carapace, PCM O FS55. O: *Bairdia* sp. D. Right lateral view of a complete carapace, PCM O FS56. P: *Bairdia* sp. E. Right lateral view of a complete carapace, PCM O FS57. Q: *Bairdia* sp. F. Right lateral view of a complete carapace, PCM O FS58. R: *Hiatabairdia subsymmetrica* Kristan-Tollmann (1970). Lateral view of a left valve, PCM O FS59.

Based on the new material this determination was revised and they are attributed to *Hungarella forelae*. Here the carapace is more triangular with a smaller radius of curvature of PB; the blade at AB is also more expressed here. The specimens described by Forel *et al.* (2019b; Plate 4, particularly fig. E, K) as *Hungarella gommerii* Forel, 2019 from the Carnian of Sichuan (South China) are very close to our specimens. However, the Sichuan specimens are smaller (biggest one  $\approx L = 600 \mu\text{m}$ ,  $H = 400 \mu\text{m}$ ) and show a smaller radius of curvature at both extremities. The largest specimen of *H. forelae* (Fig. 4) presents some morphological variability: overlap less important, at RV: the blade is located only at the

ventral part of AB and occurrence of a small spine at the upper part of it, at LV: anteroventral blade seems to be also present. We consider that these morphological variations could be the expression of ontogenic variations but some doubt remains.

*Hungarella siciliensis* n.sp.  
(Plate 1B and 1C)

**Etymology.** The species name refers to Sicily where the locus typicus is located.

**Material.** Four complete carapaces.

**Holotype.** One complete carapace, collection number PCM O 22 H 13/10/2019, Plate 1B.

**Paratype.** One complete carapace, collection number PMC O 78 P 13/10/2019, [Plate 1C](#).

**Diagnosis.** A species of *Hungarella* with triangular shape carapace, two posteroventral spines at RV, flattening in blade shape plus a spine at anterior border of RV, spine at AB of LV.

**Description.** Massive stocky carapace with a symmetric triangular shape; quite symmetric relative to H max; general shape of valves similar, but LV is significantly larger than RV and radius of curvature of PB smaller than anterior one; LV overlaps RV all around the carapace with minimum at PVB; maximum of H located at mid L or in front of mid L; maximum of L at mid H or a little below mid H; VB quite straight; presence of a very fine flattening at AB of RV in blade shape and a spine located near the maximum of convexity of AB; two more or less distinct spines at PVB of RV; one spine at AB of LV; dorsal view biconvex with valves almost symmetric in shape and W max at or just behind mid L; surface seems to be smooth.

**Dimensions.** L = 606–760  $\mu\text{m}$ ; H = 503–533  $\mu\text{m}$  (see [Fig. 4](#)).

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work).

**Remarks.** *Hungarella siciliensis* n.sp. is very close to *H. forelae* n.sp.. The shapes of the valves are similar. The main differences between the two species is the presence of 2 spines at PVB of RV, presence of a spine at AB of booth valves and the less distinct “blade” at the AB of *H. siciliensis* n.sp.. We can’t exclude that these differences are due to morphological variability of *H. forelae* n.sp. (sexual or ontogenic). However, for the time being we have not enough specimens to settle this question.

*Hungarella* sp. A  
([Plate 1D](#))

**Material.** One complete carapace.

**Dimensions.** H = 538  $\mu\text{m}$ ; L = 775  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Order Podocopida [Sars \(1866\)](#)

Suborder Bairdiocopina [Gründel \(1967\)](#)

Superfamily Bairdiodea [Sars \(1866\)](#)

Family Bairdiidae, [Sars \(1866\)](#)

Genus *Bairdia* [McCoy \(1844\)](#)

Type species *Bairdia curta* [McCoy \(1844\)](#)

*Bairdia andretrasquini* n.sp.

([Plate 1E](#) and [1F](#))

**Etymology.** In memory and honour of Dr. André Crasquin, father of the first author.

**Material.** Three complete carapaces and one broken carapace.

**Holotype.** One complete carapace, collection number PMC O 23 H 13/10/2019 ([Plate 1E](#)).

**Paratype.** One complete carapace, collection number PMC O 79 P 13/10/2019 ([Plate 1F](#)).

**Diagnosis.** A species of *Bairdia* with an elongated carapace, flattened AB and PB, and a ventral ridge along the posterior part of the VB and PB.

**Description.** Bairdioid carapace, quite elongated (H/L = 0.44); DB straight at RV and slightly convex at LV; ADB and PDB straight and quite symmetric with respect to DB; AB with large radius of curvature and maximum located

above mid-H, AB strongly flattened laterally; VB slightly concave; PB slender and pointed, maximum of curvature located at lower 1/3 of H, strongly flattened laterally; presence of the ventral ridge which begins in posterior part of VB and runs along PB.

**Remarks.** This species is quite original and differs from all other ones by the specific characters (flattened AB and PB, and a ventral ridge along the posterior part of the VB and PB.). The shape of carapace is comparable to *Bairdia* sp. 4 *sensu* [Forel \*et al.\* \(2019b\)](#) from the Carnian of China and to *Bairdia liviae* [Forel and Grădinaru \(2018\)](#) from the Anisian of North Dobrogea, Romania ([Forel and Grădinaru, 2018](#)) but this last species does not show the specific characteristics.

**Dimensions.** H = 372–415  $\mu\text{m}$ ; L = 372–415  $\mu\text{m}$ .

**Occurrence.** Tuvalian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia gambaneraensis* n.sp.

([Plate 1G](#) and [1H](#))

2018 *Bairdia* cf. *deformata* [Kollmann \(1963\)](#); Crasquin *et al.*: 134, fig. 6K-L.

**Etymology.** Referring to the locus typicus Monte Gambanera, Sicily, Italy.

**Material.** Four complete carapaces.

**Holotype.** One carapace, collection number PMC O 24H 13/10/2019 ([Plate 1G](#)).

**Paratype.** One complete carapace, collection number PMC O 80 P 13/10/2019 ([Plate 1H](#)).

**Diagnosis.** A species of *Bairdia* with a very compact carapace, a continuously arched dorsal boarder and flattened and crenulated ventral parts of AB and PB

**Description.** Bairdioid carapace, quite short (H/L = 0.6–0.7), LV overlaps RV all around the carapace with minimum at AB and anterior part of VB; LV: all the dorsal part regularly arched; AB with large radius of curvature with maximum at mid-H, VB almost straight; BP with large radius of curvature with maximum at lower 1/3 of H; PDB arched; RV: DB straight, ADB straight with an angulation of 145°–150° against DB; AB with large radius of curvature; AVB and PVB flattened laterally in its very external part and with very fine crenulation; VB slightly concave; bairdioid beak quite absent; PDB straight with an angle of 125°–130° with DB; Presence of a shoulder on medio-dorsal part of LV; carapace biconvex and quite slender in dorsal view.

**Dimensions.** L = 886–910  $\mu\text{m}$ ; H = 600–643  $\mu\text{m}$ .

**Occurrence.** Tuvalian, *Tropites dilleri* zone ([Crasquin \*et al.\*, 2018](#)), *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work).

**Remarks.** *Bairdia gambaneraensis* n.sp. is similar to *Bairdia deformata* [Kollmann \(1963\)](#) from the Rhaetian of Austria. The carapace of the present material is longer and presents a shoulder at LV. The new species has exactly the same general shape of valves as *B. penovoidea* [Bolz \(1971\)](#) from Late Norian–Rhaetian of Austria and differs from this species only by a larger AB, a longer DB and the ventral crenulation of AB and PB.

*Bairdia cassiana* ([Reuss, 1869](#))

([Plate 1I](#))

1869 *Cythere cassiana* n.sp.; Reuss: 108.

1869 *Bairdia cassiana* ([Reuss, 1869](#)); Gümbel: 180, pl. 5, figs. 18-19.

1958 *Bairdia cassiana* (Reuss, 1869); Styk: 171, fig. 3/1.  
1970 *Bairdia cassiana* (Reuss, 1869); Ulrichs: 705-706, pl. 1, figs. 1-2.

1978 *Bairdia cassiana* (Reuss, 1869); Kristan-Tollmann: 81, pl. 1, fig. 4; pl. 6, fig. 6.

1991 *Bairdia cassiana* (Reuss, 1869); Kristan-Tollmann *et al.*: 200, pl. 1, fig. 5.

1995 *Bairdia cassiana rotundidorsata* n.sp.; Monostori: 42, Pl. 2, figs. 4-5.

1996 *Bairdia (Rectobairdia) garciai* n.sp.; Crasquin-Soleau and Grădinaru: 77-78, pl. 2, figs. 5, 8.

2013 *Bairdia cassiana* (Reuss, 1869); Monostori and Tóth: 310, pl. 2, figs. 7, 8, 10.

2014 *Bairdia cassiana* (Reuss, 1869); Mette *et al.*: pl. 2, fig. 1.

2014 *Bairdia cassiana* (Reuss, 1869); Monostori and Tóth: 26, pl. 1, fig. 14.

2018 *Bairdia cassiana* (Reuss, 1869); Crasquin *et al.*: 134, fig. 6M.

2019a *Bairdia cassiana* (Reuss, 1869), Forel *et al.*, in press, figs. 4F-H.

**Material.** Four complete carapaces.

**Dimensions.** H = 486–533  $\mu\text{m}$ ; L = 840–948  $\mu\text{m}$ .

**Occurrence.** Early Carnian, Late Triassic, Southern Alps, Italy (Reuss, 1869; Gümbel, 1869; Ulrichs, 1970; Kristan-Tollmann, 1978); Carnian, Late Triassic, Święty Krzyż Mountain, Poland (Styk, 1958); Carnian, Late Triassic, Transdanubian Range, Hungary (Kristan-Tollmann, 1991); Late Anisian, Middle Triassic, Balaton Highland, Hungary (Monostori, 1995); Early Anisian, Middle Triassic, North Dobrogea, Romania (Crasquin-Soleau and Grădinaru, 1996); Ladinian, Middle Triassic, Balaton Highland, Hungary (Monostori and Tóth, 2013, 2014); Middle Anisian, Middle Triassic, Northern Calcareous Alps, Austria (Mette *et al.*, 2014); Carnian, Late Triassic, Karavanke Mountains, Slovenia (Forel *et al.*, 2019b); Tuvalian–Carnian, *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* cf. *monostorii* Forel and Grădinaru (2018)

(Plate 1J)

2018 *Bairdia* cf. *humilis* Monostori (1995); Crasquin *et al.*: 134, figs. 6i-j.

**Material.** One complete carapace and one broken carapace.

**Dimensions.** (complete carapace) H = 311  $\mu\text{m}$ ; L = 806  $\mu\text{m}$ .

**Remarks.** Forel and Grădinaru (2018) renamed *Bairdia humilis* Monostori (1995) in *Bairdia monostorii* nom. nov.

**Occurrence.** Tuvalian–Carnian, *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* sp.1 *sensu* Crasquin, Sciuto, Reitano, 2018

(Plate 1K)

2018 *Bairdia* sp. 1; Crasquin *et al.*: 134, fig. 6N-O

**Material.** One complete carapace and one right valve.

**Dimensions.** (complete carapace) H = 462  $\mu\text{m}$ ; L = 800  $\mu\text{m}$ .

**Remarks.** This species of *Bairdia* is characterized by the BD which is underlined by a blade, and by the

reticulation of the carapace. The upper part of PB is quite horizontal and its radius of curvature is small. This could be a new species.

**Occurrence.** Tuvalian–Carnian, *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* sp. A

(Plate 1L)

**Material.** Two complete carapaces.

**Dimensions.** H = 400–440  $\mu\text{m}$ ; L = 785–882  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* sp. B

(Plate 1M)

**Material.** One complete carapace.

**Dimensions.** H = 600  $\mu\text{m}$ ; L = 1000  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* sp. C

(Plate 1N)

**Material.** One complete carapace.

**Dimensions.** H = 643  $\mu\text{m}$ ; L = 1147  $\mu\text{m}$ .

**Remarks.** This species is characterized by its triangular shape, the flattening of the ventral borders and the reticulated surface. The PB has a very small radius of curvature.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* sp. D

(Plate 1O)

**Material.** One complete carapace.

**Dimensions.** H = 496  $\mu\text{m}$ ; L = 1014  $\mu\text{m}$ .

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bairdia* sp. E

(Plate 1P)

**Material.** One complete carapace.

**Dimensions.** H = 476  $\mu\text{m}$ ; L = 953  $\mu\text{m}$ .

**Remarks.** This elongated species shows a blade underlying the BD.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work).

*Bairdia* sp. F

(Plate 1Q)

**Material.** Two complete carapaces.

**Dimensions.** H = 600–615  $\mu\text{m}$ ; L = 885–953  $\mu\text{m}$ .

**Remarks.** This compact species (H/L = 0.64–0.67) has a flattened BP and AB and a shoulder at the dorsal part of the right valve.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Hiatobairdia* Kristan-Tollmann (1970)

Type species: *Hiatobairdia subsymmetrica* Kristan-Tollmann (1970)

*Hiatobairdia subsymmetrica* Kristan-Tollmann (1970) (Plate 1R)

1970 *Hiatobairdia subsymmetrica* n. gen. n.sp.; Kristan-Tollmann: 268, pl. 35, figs. 1–3.

1976 *Hiatobairdia subsymmetrica* Kristan-Tollmann (1970); Tollmann: 276, pl. 163, fig. 14.

1978 *Hiatobairdia subsymmetrica deformis* n.sp.; Kristan-Tollmann: 83, pl. 4, figs. 1–7.

1979 *Hiatobairdia subsymmetrica* Kristan-Tollmann (1970); Kristan-Tollmann *et al.*: 147, pl. 6, fig. 4.

2018 *Hiatobairdia subsymmetrica* Kristan-Tollmann (1970); Crasquin *et al.*: 134, figs. 6F–H.

**Material.** Three left valves.

**Dimensions.** H = 412–569  $\mu\text{m}$ ; L = 812–969  $\mu\text{m}$ .

**Occurrence.** Early Carnian of South Tyrol, Italy (Tollmann, 1976; Kristan-Tollmann, 1978); Rhaetian of Austrian Alps (Kristan-Tollmann, 1970) and Central Iran (Kristan-Tollmann *et al.*, 1979); Tuvanian–Carnian *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Hiatobairdia* sp. A

(Plate 2A)

**Material.** One complete carapace and one right valve.

**Dimensions.** (complete carapace) H = 533  $\mu\text{m}$ ; L = 948  $\mu\text{m}$ .

**Remarks.** This species has a straight DB and presents a ridge at the dorso-median part of the RV.

**Occurrence.** Tuvanian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Mirabairdia* Kollmann (1963)

Type species *Mirabairdia pernodosa* Kollmann (1963)

*Mirabairdia pernodosa* Kollmann (1963)

(Plate 2B)

1963 *Mirabairdia pernodosa* n.sp.; Kollmann: 177–178, pl. 1, figs. 1–2, pl. 8, figs. 1–6.

1971a *Triebelina* (*Mirabairdia*) *pernodosa illyrica* n.spp.; Kozur: 17, fig. 1G.

1971a *Triebelina* (*Mirabairdia*) *balatonica* n.sp.; Kozur: 15–16, figs. 2I, 3C.

1971 *Mirabairdia pernodosa* Kollm.; Kristan-Tollmann: text-fig. 1/8.

1984 *Triebelina* (*Mirabairdia*) *pernodosa illyrica* Kozur; Salaj and Jendrekova: pl. 2, figs. 1–4.

2014 *Triebelina* (*Mirabairdia*) *pernodosa* (Kollmann, 1963); Monostori and Tóth: 27–28, pl. 2, figs. 7–8.

**Material.** One complete carapace, 13 RV and 2 LV.

**Dimensions.** (complete carapace and LV) H (without spines) = 453–507  $\mu\text{m}$ ; L = 826–923  $\mu\text{m}$ .

**Occurrence.** Anisian, Western Carpathians, Slovakia (Salaj and Jendrekova, 1984; Kozur, 1971a); Anisian, Balaton Highland, Hungary (Kozur, 1971a); Ladinian, Dolomites, South Tirol, Italy, (Kristan-Tollmann, 1971); Ladinian, Northern Calcareous Alps, Cassian beds, Austria (Kollmann, 1963); Ladinian E-Bakony, Hungary (Monostori and Tóth, 2014); Tuvanian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Ptychobairdia* Kollmann (1960)

Type species *Ptychobairdia kuepperi* Kollmann (1960)

*Ptychobairdia iudicaensis* n.sp.

(Plate 2C and 2D)

**Etymology.** From the locus typicus Castel di Iudica, Sicily, Italy.

**Material.** Four complete carapaces.

**Holotype.** One complete carapace, collection number PMC O 25 H 13/10/2019 (Plate 2C).

**Paratype.** One complete carapace, collection number PMC O 81 P 13/10/2019 (Plate 2D).

**Diagnosis.** A species of *Ptychobairdia* with a reticulated carapace which is flattened laterally all around except at the ventral part; LV significantly higher than RV, presence of vertical sulci at antero-dorsal part of the carapace.

**Description.** Carapace massive, high (H/L = 0.6); surface reticulated;

LV: Flattened laterally all around with maximum at DB and PB; BD strongly arched; AB with quite large radius of curvature and maximum at mid H; VB almost straight; BP with a very small curvature; two vertical sulci in dorsal part; LV strongly overlapping RV all around with maximum at BD.

RV: Strongly flattened all around except in ventral part; presence of a sulcus in AD part; BD long; AB with quite small radius of curvature; VB gently concave at its anterior part; BP very slender; DB, ADB, AVB, PVB, PDB straight.

**Remarks.** *Ptychobairdia iudicaensis* n.sp. is comparable with *P. oberhauseri* Kollmann (1960) from the Rhaetian of Austria (Kollmann, 1960) and the Carnian–Norian of Queen Charlotte Island, Canada (Arias and Lord, 2000). The latter species is longer, has a smaller AB and shows a horizontal sulcus. *P. iudicaensis* n.sp. differs from *P. kristanae* Kollmann (1960) from the Rhaetian–Early Jurassic of Austria (Kollmann, 1960, 1963) and the late Carnian of Sicily (Crasquin *et al.*, 2018) by its reticulated carapace and the RV being clearly smaller than LV.

**Dimensions.** L = 720–1083  $\mu\text{m}$ ; H = 480–667  $\mu\text{m}$  (see Fig. 5).

**Occurrence.** Tuvanian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Ptychobairdia leonardo* n.sp.

(Plate 2E and 2F)

**Etymology.** Dedicated to Leonardo Reitano, son of Agatino Reitano.


**Material.** Six right valves, eight left valves.

**Holotype.** One right valve, collection number PMC O 26 H 13/10/2019 (Plate 2E).

**Paratype.** One left valve, collection number PMC O 82 P 13/10/2019 (Plate 2F).

**Diagnosis.** A species of *Ptychobairdia* with a short coarse reticulated carapace, strongly compressed and finely reticulated AB and PB and a central node.

**Description.** Carapace with massive coarse reticulation, flattened laterally at AB and PB; DB straight at both valves and parallel to VB; ontogenic modifications of DB: at RV with nodules or blade at biggest specimens, at LV development of shoulders at each extremities in largest specimens; AB and PB with small radius of curvature, flattened laterally and covered by a fine reticulation; VB straight to slightly concave, with



**Plate 2.** Ostracods from Late Triassic (Tuvalian-Carnian) of Monte Gambanera, Sicily, Italy. All the specimens are stored in the Palaeontological Museum of the University of Catania. The repository numbers are given as PMC (Palaeontological Museum Catania) O (Ostracods) X H (number of holotype) or X P (number of paratype) or FS X (Figured Specimen number) registration date. Scale bars = 200  $\mu\text{m}$  except P-Q, R = 100  $\mu\text{m}$ . A: *Hiatobairdia* sp. A. Right lateral view of a complete carapace, PMC O FS61. B: *Mirabairdia pernodosa* Tollmann, 1963. Right lateral view of a complete carapace, PMC O FS61. C-D: *Ptychobairdia iudicaensis* n.sp. C: holotype, right lateral view of a complete carapace, PMC O25 H 13/10/2019; D: paratype, right lateral view of a complete carapace, PMC O 81 P 13/10/2019. E-F: *Ptychobairdia leonardo* n.sp. E: holotype, lateral view of a right valve, PMC O 26 H 13/10/2019; F: paratype, lateral view of a left valve, PMC O 82 P 13/10/2019. G-H: *Petasobairdia jeandercourti* n.sp. G: holotype, right lateral view of a complete carapace, PMC O 27 H 13/10/2019; H: paratype, left lateral view of a complete carapace, PMC O 83 P 13/10/2019. I: *Urobairdia angusta* Kollmann (1963). Right lateral view of a complete carapace, PCM O FS62. J: *Bairdiacypris triassica* Kozur (1971c). Right lateral view of a complete carapace, PCM O FS63. K: *Bythocypris* sp. A. Left lateral view of a complete carapace, PCM O FS64. L: *Acratia maugerii* Crasquin *et al.* (2018). Left lateral view of a complete carapace, PCM O FS65. M-N: *Kerocythere dittainoensis* n.sp. M: holotype, lateral view of a right valve, PMC O 28 H 13/10/2019; N: paratype, lateral view of a left valve, PMC O 84 P 13/10/2019. O: *Renngartenella sanctaeruscis* Kristan-Tollmann (1973) ( $\delta$ ). Right lateral view of a complete carapace, PCM O FS66. P-Q: *Mockella barbroyae* n.sp. P: holotype, right lateral view of a complete carapace, PMC O 29 H 13/10/2019; Q: paratype, right lateral view of a complete carapace, number PMC O 85 P 13/10/2019. R: *Simeonella brotzenorum* Sohn (1968). Lateral view of a right valve, PCM O FS67.

development of adventral structure; presence of a big node in median part of the carapace.

**Remarks.** *Ptychobairdia leonardo* n.sp. differs from other species by the specific characters. *Margarobairdia zapfei* Kristan-Tollmann (1983) from the Anisian of South China (Kristan-Tollmann, 1983) has a similar valve shape but a different ornamentation.

**Dimensions.** L = 610–776  $\mu\text{m}$ ; H = 362–553  $\mu\text{m}$  (see Fig. 6).

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Petasobairdia* Chen and Shi (1982)

Type species *Petasobairdia bicornuta* Chen and Shi (1982)

*Petasobairdia jeandercourti* n.sp.  
(Plate 2G and 2H)

**Etymology.** Dedicated to past Pr. Jean Dercourt, who was the mentor of the first author.


**Material.** One broken carapace and three left valves.

**Holotype.** One broken carapace, collection number PMC O 27 H 13/10/2019 (Plate 2G).

**Paratype.** One left valve, collection number PMC O 83 P 13/10/2019 (Plate 2H).

**Diagnosis.** A species of *Petasobairdia* with a long reticulated carapace and elongated nodes at ADB and PDB of both valves.

**Description.** Carapace long (H/L  $\approx$  0.4), reticulated, strongly laterally compressed along AB, AVB, VB, PVB,



**Fig. 4.** Height (H)/length (L) diagram of figured specimens of the two new *Hungarella* species. In blue: *H. forelae* n.sp.; in orange: *H. siciliensis* n.sp.

PB; DB long and straight at both valves; presence of an elongated node at ADB and PDB of both valves; LV with two big horns with large base at each extremities of DB.

**Dimensions.** (holotype and paratype without spines) H = 348–373 µm; L = 826–853 µm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** *Ptychobairdia jeandercourti* n.sp. is comparable to *P. bicornuta* Chen and Shi (1982) from the Late Permian of Hubei Province (Chen and Shi, 1982) which has a much shorter DB. *P. longispinosa* (Kozur, 1971a, b, c) from Anisian of Slovakia (Kozur, 1971a), Anisian and Middle Triassic of Romania (Salaj and Jendrekova, 1984; Crasquin-Soleau and Grădinaru, 1996; Sebe *et al.*, 2013), Anisian of Austria (Mette *et al.*, 2014), Ladinian of Hungary (Monostori and Tóth, 2013) and Carnian of Southern Turkey (Forel *et al.*, 2017) has a shorter DB and doesn't have AD and PD nodes.

Genus *Urobairdia* Kollmann (1963)

Type species: *Urobairdia austriaca* Kollmann (1963)

*Urobairdia angusta* Kollmann (1963)

(Plate 2I)

1963 *Urobairdia angusta* n.g. n.sp.; Kollmann: 167, pl. 6, figs. 1–4.

1965 *Urobairdia angusta* Kollmann (1963); Széles: 414, pl. 6, fig. 5.

1995 *Bairdia (Urobairdia) angusta recta* n.sp.; Monostori: 43, pl. 3, figs. 3–4.

2010 *Urobairdia angusta* Kollmann (1963); Zorn: 271–272, pl. 6, figs. 12–15.


2013 *Bairdia (Urobairdia) angusta* Kollmann (1963); Monostori and Tóth: 7, pl. 1, figs. 10–12.

2014 *Bairdia (Urobairdia) angusta* Kollmann (1963); Monostori and Tóth: 25–26, Pl. 1, fig. 14.


**Material.** Two complete carapaces

**Dimensions.** H = 440–500 µm; L = 826–871 µm.

**Occurrence.** Late Norian, Zlambach Formation, Austria (Kollmann, 1963; Zorn, 2010); Anisian, Felsőörs, Hungary (Monostori, 1995); Carnian, Nosztori Valley, Hungary (Széles, 1965); Ladinian–Carnian, Balaton Highland, Hungary (Monostori and Tóth, 2013, 2014); Tuvalian–Carnian,



**Fig. 5.** Height (H)/length (L) diagram for *Ptychobairdia iudicaensis* n.sp. In red: right valves; in blue left valves.



**Fig. 6.** Height (H)/length (L) diagram for *Ptychobairdia leonardoii* n.sp. In blue: left valves; in red: right valves.

*Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Genus *Bairdiacypris* Bradfield (1935)

Type species: *Bairdiacypris deloi* Bradfield (1935).

*Bairdiacypris triassica* Kozur (1971a, b, c)

(Plate 2J)

1911 ?*Bairdia silicula* Jones; Méhés: 16–17, pl. 1, figs. 19–21

1971 *Bairdiacypris triassica* n.sp.; Kozur: 5–6, figs. 2H–L

2013 *Bairdiacypris triassica* Kozur (1971a, b, c); Monostori and Tóth: 313–314, pl. 3, figs. 7–8

2014 *Bairdiacypris triassica* Kozur (1971a, b, c); Monostori and Tóth: 25, pl. 1, fig. 12

2017 *Bairdiacypris triassica* Kozur (1971c); Forel *et al.*: fig. 10U

**Material.** Two complete carapaces

**Dimensions.** H = 361–374 µm; L = 774–812 µm

**Occurrence.** Early Carnian, Balaton highland, Hungary (Méhés, 1911; Kozur, 1971c), Ladinian, Nosztori Valley, Hungary (Monostori and Tóth, 2013); Ladinian–Carnian, Balaton Highland (Monostori and Tóth, 2014), Carnian, Mersin, Turkey (Forel *et al.*, 2017); Tuvalian–Carnian,

*Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Family Bythocypridae Maddocks (1969)

Genus *Bythocypris* Brady (1880)

Type species: *Bythocypris reniformis* Brady (1880)

*Bythocypris* sp. A

(Plate 2K)

**Material.** Two complete carapaces and two LV.

**Dimensions.** H = 433–500 µm; L = 775–1090 µm.

**Occurrence.** Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Family Acratiidae Gründel (1962)

Genus *Acratia* Delo (1930)

Type species: *Acratia typica* Delo (1930).

*Acratia maugerii* Crasquin *et al.* (2018)

(Plate 2L)

1991 *Acratia* sp.; Kristan-Tollmann: 196, pl. 1, fig. 1.

2013 *Acratia goemoeryi* (Kozur, 1970); Monostori and Tóth: 6–7, pl. 4, only fig. 2.

2018 *Acratia maugerii* n.sp.; Crasquin *et al.*: 137–138, fig. 7H–J.

**Material.** One complete carapace and two LV.

**Dimensions.** H = 373–464 µm; L = 866–1066 µm.

**Occurrence.** Late Ladinian of NE Iran (Kristan-Tollmann, 1991), Balaton Highland, Hungary (Monostori and Tóth, 2013); Tuvalian–Carnian, *Tropites dilleri* zone (Crasquin *et al.*, 2018) and *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Superfamily Cytheroidea Baird (1850)

Family Cytheruridae Müller (1894)

Subfamily Cytherurinae Müller (1894)

Genus *Kerocythere* Kozur and Nicklas (1970)

Type species *Cythere raibliana* Gümbel (1869)

*Kerocythere dittainoensis* n.sp.

(Plate 2M and 2N)

**Etymology.** Referring to the Dittaino river which flows near to the locus typicus.

**Material.** Four complete carapaces.

**Holotype.** One complete carapace, collection number PMC O 28 H 13/10/2019 (Plate 2M).

**Paratype.** One complete carapace, collection number PMC O 84 P 13/10/2019 (Plate 2N).

**Diagnosis.** A species of *Kerocythere* Kozur and Nicklas (1970) with a subrectangular reticulate carapace, presence of a lateral thick ridge which ascends at PB and occurrence of ventral ridges, one thick and several thinner ones parallel to VB.

**Description.** Carapace subrectangular, almost equilateral; BD long and straight, presence of a ridge on each side of hinge; presence of an eye spot; AB with large radius of curvature with maximum located below mid-H, flattened laterally and smooth; VB almost straight; PB with small radius of curvature with maximum around mid H, upper and lower part quite straight; H max at anterior angle; L max at PB; sulcus more or less developed in anterior 1/3 of L; surface reticulated and ornamented with possible pustules and ridges: one lateral,

thick, reaching from antero-ventral part of the carapace up to PB, ascending in posterior part; group of ventral ridges, one thick parallel to VB and several (at least three) below.

**Dimensions.** L = 651–731 µm; H = 309–376 µm.

**Occurrence.** *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** *Kerocythere dittainoensis* n.sp. could be compared to *Kerocythere reticulata* Kristan-Tollmann (1972) from early Carnian of the Dolomites (Italy). This species doesn't show the ventral group of ridges but has one ridge at the AD part of the carapace following the AB. The new species is also close to *Kerocythere tricostata* Forel, 2017 from the middle Carnian of southern Tauride-Anatolide platform (Turkey; Forel *et al.*, 2017). At the present material the lateral ridge is longer, ascends at its posterior part and the surface is reticulated.

Family indet.

Genus *Renngartenella* Schneider 1957 (in Mandelstam *et al.*, 1957)

*Renngartenella sanctaecrucis* Kristan-Tollmann (1973)

(Plate 2O)

1973 *Renngartenella sanctaecrucis* Kristan-Tollmann; Kristan-Tollmann and Hamedani: 215, 217–219, pl. 8, figs. 1–6; pl. 11, figs. 1, 3, 5, 6; pl. 12, fig. 10.

1979 *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Lieberman: 215, pl. 5, fig. 2.

1982 *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Basha: pl. 1, fig. 15.

1990 *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Gerry *et al.*: 96, pl. 1, figs 11–13.

1994 *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Monostori: 320, 322, figs 5/5–7.

2001 *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Keim *et al.*: fig. 8C.

2014 *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Monostori and Tóth: 29–30, pl.3, figs. 10–12.

2019a *Renngartenella sanctaecrucis* Kristan-Tollmann (1973); Forel *et al.*: fig. 7T–U, in press.

**Material.** Ten carapaces.

**Dimensions.** H = 316–439 µm; L = 567–900 µm.

**Occurrence.** Julian, early Carnian, Heiligkreuz Formation, Italy (Kristan-Tollmann and Hamedani, 1973); Carnian, Late Triassic, Italian Alps (Lieberman, 1979); Cordevolian, Carnian, Jordan (Basha, 1982); Carnian, Israel (Gerry *et al.*, 1990); Carnian, Balaton Highland, Hungary (Monostori 1994; Monostori and Tóth, 2014); Carnian, Dolomites, Northern Italy (Keim *et al.*, 2001); Carnian, Karavanke Mountains, Slovenia (Forel *et al.*, 2019b); Tuvalian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Superfamily Cypridoidea Baird (1845)


Family Limnocytheridae Klie (1938)

Genus *Mockella* Bunza and Kozur (1971)

Type species: *Mockella muelleri* Bunza and Kozur (1971); subsequent designation (Kozur, 1973)

*Mockella barbroae* n.sp.

(Plate 2P and 2Q)



**Fig. 7.** Height (H)/length (L) diagram for *Mockella barbroae* n. sp.

2018 *Mockella muelleri* Bunza and Kozur (1971), Crasquin *et al.*: 139, figs. 70-P.

**Etymology.** Personal dedication of the first author to Mrs. Barbro Lamy, in token of friendship and affection.

**Material.** Seven complete carapaces and two carapaces from Crasquin *et al.* (2018)

**Holotype.** One complete carapace, collection number PMC O 29 H 13/10/2019 (Plate 2P).

**Paratype.** One complete carapace, collection number PMC O 85 P 13/10/2019 (Plate 2Q).

**Diagnosis.** A species of *Mockella* with a long subrectangular carapace and a well-developed rib all around the carapace.

**Description.** Carapace sub rectangular, long ( $H/L = 0.535$ ); eye spot well developed; AB with maximum of curvature located low between mid-H and lower 1/3 of H; VB straight; PB equivalent to AB in heteromorphs and smaller in tecnomorphs with maximum of curvature located above mid-H; anteromedian sulcus located in front of mid-L; posterior lobe well developed; anterior lobe less distinct; presence of a distinct ridge all around the carapace including BD; presence of additional ridges on lateral surface of the valves:

- a median ridge which begins at maximum of curvature of AB and precedes on the posterior lobe; this ridge is high and stands out on the surface;
- a lateral ridge below the lobes parallel to VB;
- a small ridge in upper part of AB and below the eye spot.

The valve surface is reticulated with 4 small pustules distributed parallel to AB; in dorsal view, the flanks are parallel. Sexual dimorphism present, expressed by the thickness of the posterior part of the carapace in heteromorphs.

**Dimensions.**  $L = 270\text{--}492\ \mu\text{m}$ ;  $H = 150\text{--}275\ \mu\text{m}$  (see Fig. 7).

**Occurrence.** Late Carnian (*Tropites dilleri* zone), Mufara Formation, Sicily, Italy Tuvallian–Carnian (Crasquin *et al.*, 2018) and Tuvallian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

**Remarks.** *Mockella barbroae* n.sp. is very close to *M. muelleri* Bunza and Kozur (1971) from the late Carnian of Tyrol, Austria (Bunza and Kozur, 1971) and the Carnian of

Monte Cammarata, Sicily (Cafiero and De Capoa Bonardi, 1982). In the study on the Mufara Formation (Crasquin *et al.*, 2018) we attributed the specimens to the latter species. The very well preserved material enabled us to review our attribution. At the present specimens the BP is larger, the median ridge ends at the posterior lobe and doesn't reach the BP; an additional ridge is present below the lobes and the flanks are parallel in dorsal view.

Genus *Simeonella* Sohn (1968)

Type species: *Simeonella brotzenorum* Sohn (1968).

*Simeonella brotzenorum* Sohn (1968)

(Plate 2R)

1968 *Simeonella brotzenorum* n.sp.; Sohn: 23-24, pl. 2, figs. 1-4, 6-8, 12-22.

1971 *Simeonella brotzenorum alpina* n.sp.; Bunza and Kozur: 4-5, pl. 1, figs. 5-7, 13.

1971 *Simeonella brotzenorum norica* n.sp.; Bunza and Kozur: 5-6, pl. 1, fig. 3.

1973 *Simeonella brotzenorum* Sohn (1968); Kristan-Tollmann and Hamedani: text-fig. 13/2.

1974 *Simeonella brotzenorum* Sohn (1968); Hirsch and Gerry: pl. 2, figs. 1-2.

1979 *Simeonella brotzenorum* Sohn (1968); Lieberman: 103, pl. 5, figs. 6-7.

1979 *Simeonella brotzenorum alpina* Bunza and Kozur (1971); Styk: 119, pl. 28, figs. 9-10.

1982 *Simeonella brotzenorum* Sohn (1968); Basha: pl. 1, fig. 11.

1990 *Simeonella brotzenorum* Sohn (1968); Gerry *et al.*: 95, pl. 1, figs. 3-5.

1994 *Simeonella brotzenorum nostorica* n.sp.; Monostori: 324-325, text-fig. 6/1-6.

2001 *Simeonella brotzenorum nostorica* Monostori (1994); Keim *et al.*: fig. 8B.

**Material.** Two carapaces.

**Dimensions.**  $H = 269\text{--}296\ \mu\text{m}$ ;  $L = 446\text{--}488\ \mu\text{m}$ .

**Occurrence.** Ladinian to Carnian, Makhtesh Ramon, Israel (Sohn, 1968; Hirsch and Gerry, 1974; Gerry *et al.*, 1990); Carnian, Northern Calcareous Alps, Austria (Bunza and Kozur, 1971; Kristan-Tollmann and Hamedani, 1973); Carnian, Julian Alps, Italy (Lieberman, 1979; Keim *et al.*, 2001); Carnian, Poland (Styk, 1958), Carnian, Jordan Valley, Jordan (Basha, 1982); Carnian, Balaton Highland, Hungary (Monostori, 1994; Monostori and Tóth, 2014); Tuvallian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this work).

Family Candonidae Kaufmann (1900)

Genus *Paracypris* Sars (1866)

*Paracypris?* cf. *redcarensis* (Blake, 1876)

(Plate 3A)

**Material.** One complete carapace.

**Dimensions.**  $H = 257\ \mu\text{m}$ ;  $L = 561\ \mu\text{m}$ .


**Occurrence.** Tuvallian–Carnian, *Tropites subbullatus/Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Paracypris?* sp. A

(Plate 3B)

**Material.** One carapace.

**Dimensions.**  $H = 442\ \mu\text{m}$ ;  $L = 631\ \mu\text{m}$ .



**Plate 3.** Ostracods from Late Triassic (Tuvlian-Carnian) of Monte Gambanera, Sicily, Italy. All the specimens are stored in the Palaeontological Museum of the University of Catania. The repository numbers are given as PCM (Palaeontological Museum Catania) O (Ostracods) FS X (Figured Specimen number) registration date. A, B, C, D: Scale bars = 200  $\mu\text{m}$ ; E, F, G, H, I: scale bars = 100  $\mu\text{m}$ . A: *Paracypris?* cf. *redcarensis* (Blake, 1876). Lateral view of a right valve, PCM O FS68. B: *Paracypris?* sp. A. Right lateral view of a complete carapace, PCM O FS68. C: *Bektasia* sp. A. Right lateral view of a complete carapace, PCM O FS69. D: *Bektasia* sp. B. Right lateral view of a complete carapace, PCM O FS70. E: Podocopida gen. sp. indet. Lateral view of a right valve, PCM O FS71. F: *Polycope baudi* Crasquin-Soleau and Grădinaru (1996). Lateral view of a complete carapace, PCM O FS72. G: *Polycope densoreticulata* Monostori and Tóth (2013). Lateral view of a complete carapace, PCM O FS73. H: *Polycope* sp. A. Lateral view of a complete carapace, PCM O FS74. I: *Thaumatomma?* sp. A. Right lateral view of a complete carapace, PCM O FS75.

**Occurrence.** Tuvlian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Order Platycopida Sars (1866)  
 Suborder Platycopina Sars (1866)  
 Superfamily Cavellinoidea Egorov (1950)  
 Family Cavellinidae Egorov (1950)  
 Genus *Bektasia* Özdikmen (2010)  
 Type species: *Reubenella avnimelechi* Sohn (1968).  
*Bektasia* sp. A  
 (Plate 3C)

**Material.** One carapace.

**Dimensions.** H = 477  $\mu\text{m}$ ; L = 787  $\mu\text{m}$ .

**Occurrence.** Tuvlian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Bektasia* sp. B  
 (Plate 3D)

**Material.** One carapace.

**Dimensions.** H = 545  $\mu\text{m}$ ; L = 763  $\mu\text{m}$ .

**Occurrence.** Tuvlian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Order, genus and species indet.

Podocopida gen. sp. indet.

(Plate 3E)

**Material.** One carapace.

**Dimensions.** H = 188  $\mu\text{m}$ ; L = 364  $\mu\text{m}$ .

**Occurrence.** Tuvlian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Subclass Myodocopa Sars (1866)  
 Order Myodocopida Sars (1866)  
 Superfamily Polycopidea Sars (1866)  
 Family Polycopidae Sars (1866)  
 Genus *Polycope* Sars (1866)  
*Polycope baudi* Crasquin-Soleau and Grădinaru (1996)  
 (Plate 3F)

1996 *Polycope baudi* n.sp.; Crasquin and Grădinaru: 15–16, figs. 1–3.

2013 *Polycope baudi* Crasquin and Grădinaru 1996; Sebe *et al.*: pl. 1, fig. 1.

**Material.** Two carapaces.

**Dimensions.** H = 330–328  $\mu\text{m}$ ; L = 357–376  $\mu\text{m}$ .

**Occurrences.** Early–middle Anisian, Uzum Bair, Dobrogea, Roumania (Crasquin-Soleau and Grădinaru, 1996; Sebe *et al.*, 2013); Tuvlian–Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).


?*Polycope densoreticulata* Monostori and Tóth (2013)  
 (Plate 3G)

2013 *Polycope densoreticulata* n.sp.; Monostori and Tóth: 5, pl. 1, figs. 4–7.

**Material.** Three carapaces.

**Dimensions.** H = 204–293  $\mu\text{m}$ ; L = 231–306  $\mu\text{m}$ .

**Occurrence.** Ladinian, Balaton Highland, Hungary (Monostori and Tóth, 2013); Tuvlian–Carnian, *Tropites*



**Fig. 8.** Diversity of ostracod families from the *Tropites subbullatus*/*Anatropites spinosus* zones represented by number of genera (A) and species (B) in the samples of Mount Gambanera.

*subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

*Polycope* sp. A  
(Plate 3H)

**Material.** One carapace.

**Dimensions.** H = 289 µm; L = 278 µm.

**Occurrences.** *Tuvalian–Carnian*, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).

Order Halocyprida Dana (1853)

Suborder Halocypridina Dana (1853)

Superfamily Thaumatoocypridoidea Müller (1906)

Family Thaumatoocyprididae Müller (1906)

Genus *Thaumatomma* Kornicker and Sohn (1976)

Type species: *Thaumatomma piscifrons* Kornicker and Sohn (1976)


*Thaumatomma* ? sp.A

(Plate 3I)

**Material.** Two carapaces.

**Dimensions.** H = 525–600 µm; L = 575–600 µm.

**Occurrence.** *Tuvalian–Carnian*, *Tropites subbullatus*/*Anatropites spinosus* zones, Monte Gambanera, Central-Eastern Sicily, Italy (this study).




**Fig. 9.** Diversity of ostracod families from the *Tropites dilleri* zone represented by the number of genera (A) and species (B) in the samples of Mount Scalpello (data from Crasquin *et al.*, 2018).

## 6 Assemblage analysis

The assemblage is composed of 200 specimens belonging to 10 families (plus two undetermined families), 19 genera and 37 species. The 10 determined families present are: Healdiidae, Bairdiidae, Bythocyprididae, Acratidae, Cytheruridae, Limnocytheridae, Candonidae, Cavellinidae, Polycopidae, Thaumatoocyprididae. The relative abundance of the different families expressed by the numbers of genera and species is given in Figure 8.

The Bairdiidae, the most abundant family in number of species (53%) and genera (37%) (Fig. 8), are present in marine environments ranging from very shallow waters up to deep seas. The morphology of the family changes from massive thick-shelled forms in nearshore environments to elongate thin-shelled forms beyond continental slope (particularly in genus *Bairdia*). In the same way, the carapaces of Acratidae lengthen with depth (as example: *Acratia goemeryi* Kozur (1970) from Early-Smithian- to Late-Carnian-Triassic; see Forel *et al.*, 2017). Here these two families present thick and ornamented shells (Plates 1E–1R and 2A–2L) which testify an open marine environment with moderate energy. The Healdiidae do also show a change of morphology with depth. The specimens with massive shells and small spines are neritic



**Fig. 10.** Palaeogeographic reconstruction of Tethyan (left) and central Mediterranean (right) areas during Late Triassic (after Di Stefano *et al.*, 2015, modified).

inhabitants (Plate 1A–1D) of relatively nearshore muddy conditions. In the deep sea, the specimens are thin shelled, elongate with long spines (*e.g.* Palaeozoic genus *Timorhealdia* Bless, 1987). In other families, some genera also show different morphological adaptation from neritic to deep sea environments (*Healdia*, *Microcheilinella* etc. see synthesis Tab. 1 in Crasquin and Horne). The ratios between these different morphologies have been used to characterize the depth of late Palaeozoic and Early Mesozoic environments since quite a long time (model of Lethiers and Raymond, 1991). The present assemblage doesn't include any unequivocal deep water taxa such as those discovered recently in the Carnian of Southern Turkey for example (Forel *et al.*, 2017) or of Sichuan, South China (Forel *et al.*, 2019b). Although the number of specimens is very low, the diversity is quite high with 10 determined families (plus 2 undetermined), 17 genera and 37 species. This biodiversity testifies normal marine conditions and absence of environmental stress. The taxon *Simeonella brotzenorum* Sohn (1968) which is characteristic of brackish–hypersaline conditions (Gerry *et al.*, 1990; Monostori, 1994) is present but with only 2 carapaces. Quite all the specimens are preserved with the complete carapace. After the death of the specimens, the carapaces tend to open in a few hours (Guernet and Lethiers, 1989). This could suggest a rapid burial in the sediments due to a high sedimentation rate. Similar taphonomic characteristics were also found by Pokorný (1964) and Oertli (1971) for pelitic layer associations deposited in basins with extremely rapid distal sedimentation. We compare the results of the *Tropites subbullatus/Anatropites spinosus* zones (this study), with the data obtained in levels just below in *Tropites dilleri* zone (Crasquin *et al.*, 2018) (Fig. 9). Through time, the assemblage became more diversified as recorded by the increasing number of families (8 to 12), genera (14 to 18) and species (23 to 36). The percentage of Bairdiidae decreases in favour of two families being absent before, Cytheruridae and Limnocytheridae which include typical genera of nearshore environments such as *Simeonella*, *Mockella* and *Kerocythere*. We observe also the presence

of the brackish–hypersaline species *Renngartenella sanctaecrusis* Kristan-Tolmann, which was suggested by Gerry *et al.* (1990) to be a stenohaline ostracod.

For Monostori (1994), the dominance of three genera *Kerocythere*–*Renngartenella*–*Simeonella* seems to be a signal of salinity variability. Although these genera are not dominant here, their presence testifies a shallowing of environment from the *Tropites dilleri* zone to the *Tropites subbullatus/Anatropites spinosus* zones. The genus *Acratia* is a typical Palaeozoic form present both in Eifelian (neritic) and Thuringian (deep) mega-assemblages (see synthesis in Crasquin and Horne, 2018). Nevertheless, this genus survived the Permian–Triassic extinction events. In 2013, Crasquin and Forel mentioned the last occurrence of neritic *Acratia* in the Spathian and of deep marine *Acratia* in the Anisian (Crasquin-Soleau and Grădinaru, 1996). Since then, some deep marine forms were also found in the Ladinian of Balaton Highland (Monostori and Tóth, 2013), in the Carnian of Turkey (Forel *et al.*, 2017) and Slovenia (Forel *et al.*, 2019b). In 2013, Monostori and Tóth, described *Acratia goemeryi* from Ladinian neritic sediments of a borehole in Hungary. Three species of *Acratia* from the Carnian of Karavanke Mountains, Solvenia, are figured in Forel *et al.* (2019b). The occurrence of *Acratia maugerii* in the present material confirms that *Acratia* occurs in neritic environments of the Carnian.

## 7 Conclusion

The palaeoecological interpretation of the sedimentary facies of the Mufara Formation is extremely difficult due to the absence of intact outcrops. The original stratification and the sedimentary structures have been completely destroyed because of continuous agricultural processing of the pelitic soils and their very consistency which determines frequent drifts and landslides. In very few and limited locations parallel laminations and sandy levels were observed. The only useful palaeoecologic data are those obtained from the




Fig. 11. Schematic palaeoenvironmental model for the late Carnian Mufara Formation basin (see also Fig. 10).

palaeontological analysis. The results of the ostracod fauna analysis allow the following conclusions:

- 1 The ostracod assemblage doesn't yield any evidence of deep marine taxa both at Mt. Scalpello (Crasquin *et al.*, 2018) and at Mt. Gambanera. The Mufara Basin, therefore, can be interpreted as a shallow marine basin (Fig. 10) within the deepest and most distal part of a vast continental shelf where the carbonate platforms Panormide, Trapanese, Saccense were located. The deep marine ostracod fauna discovered recently in the Carnian of Southern Turkey (Forel *et al.*, 2017) or in the South China (Forel *et al.*, 2019a) suggests a deepening of the Neo-Tethys basin towards the more eastern areas.
- 2 The Mufara Basin was a site of rapid and intense sedimentation probably linked to rapid bottom currents which, sometimes, displaced and transported (also *in vivo*) microfaunas from more superficial neighbouring environments. It is pointed out here that the sediments of the Mufara Basin at Monte Gambanera do not show vortex structures which were recognized in the Mufara Basin at Monte Scapello. This suggests, that the sediment environment of the Mufara Formation outcrops at Monte Gambanera (Fig. 11) was more distal and less turbulent than that of Mt which was effected by strong bottom traction and swirling currents (Crasquin *et al.*, 2018).
- 3 According to many authors, the Mufara basin is located in a transitional position between the bathyal Neotethys facies to the south and southeast and the carbonate platforms that surround it (Figs. 10 and 11).

### Taxa registration:

*Hungarella forelae*: [urn:lsid:zoobank.org:act:DE0CE7FE-10E0-4F8E-8DC8-C829D3D5485B](https://zoobank.org/act:DE0CE7FE-10E0-4F8E-8DC8-C829D3D5485B)

*Hungarella siciliensis*: [urn:lsid:zoobank.org:act:942B09CB-1014-4CD3-A244-4EC52F0633B9](https://zoobank.org/act:942B09CB-1014-4CD3-A244-4EC52F0633B9)

*Bairdia andreocrasquini*: [urn:lsid:zoobank.org:act:B42972B5-54DF-4435-9C2B-E3DD46610140](https://zoobank.org/act:B42972B5-54DF-4435-9C2B-E3DD46610140)

*Bairdia gambaneraensis*: [urn:lsid:zoobank.org:act:DAB3F723-F5D1-40B1-B6BD-CC37BC92DD82](https://zoobank.org/act:DAB3F723-F5D1-40B1-B6BD-CC37BC92DD82)

*Ptychobairdia iudicaensis*: [urn:lsid:zoobank.org:act:FF725BE-043C-4295-AD53-9023B9321380](https://zoobank.org/act:FF725BE-043C-4295-AD53-9023B9321380)

*Ptychobairdia leonardoii*: [urn:lsid:zoobank.org:act:FC93D70B-B0C0-4898-85BC-A4F3DA21E560](https://zoobank.org/act:FC93D70B-B0C0-4898-85BC-A4F3DA21E560)

*Petasobairdia jeandercourti*: [urn:lsid:zoobank.org:act:84DA8AAD-D794-4F58-A432-531D6DE12EBF](https://zoobank.org/act:84DA8AAD-D794-4F58-A432-531D6DE12EBF)

*Kerocythere dittainoensis*: [urn:lsid:zoobank.org:act:E47E2789-5811-4B0E-B05C-9C5E6AAC6216](https://zoobank.org/act:E47E2789-5811-4B0E-B05C-9C5E6AAC6216)

*Mockella barbroae*: [urn:lsid:zoobank.org:act:4ADD818E-6B13-4162-B348-1A807B0CF100](https://zoobank.org/act:4ADD818E-6B13-4162-B348-1A807B0CF100)

*Acknowledgements.* The authors are grateful to the reviewers and the editors for detailed suggestions and comments to the manuscript. Thanks are due to Mr. Alfio Viola (Electronic Microscopy Laboratory, University of Catania) for SEM photos assistance. Catania Palaeoecological Research Group contribution n°456.

We thanks the two reviewers Emőke Tóth from Eötvös Loránd University, Budapest, Hungary and Wolfgang Mette from Innsbruck University, Austria for their fruitful comments to improve our paper.

Palaeoecological Research Group contribution n°456. This research was supported by the University of Catania “Piano della Ricerca 2016/2018 (code no. 227221132118)” and “PiaCeRi–Piano Incentivi per la Ricerca di Ateneo 2020-22 linea di intervento 2”.

### References

- Anderson FW. 1964. Rhaetic Ostracoda. *Bulletin of the Geological Survey of Great Britain* 21: 133–174.
- Arias C, Lord A. 2000. Upper Triassic marine Ostracoda from the Queen Charlotte Islands British Columbia Canada. *Revista Española de Micropaleontología* 32: 175–192.
- Baird W. 1845. Arrangement of the British Entomostraca, with a list of species, particularly noticing those which have as yet been discovered within the bounds of the club. *Transactions of Berwick's Naturalist Club* 2: 145–158.
- Baird W. 1850. The Natural History of the British Entomostraca. London: Ray Society, 364 p.
- Basha SHS. 1982. Microfauna from the Triassic rocks of Jordan. *Revue de Micropaléontologie* 25: 3–11.
- Blake JE. 1876. Subclass Entomostraca. In: Late R, Blake JR, eds. *The Yorkshire Lias*, pp. 429–435.
- Bless MJM. 1987. Lower Permian ostracodes from Timor (Indonesia). *Proceedings of Koninklijke Nederlandse Akademie van Wetenschappen* 90: 1–13.
- Bolz H. 1971. Die Zlambach-Schichten (alpine Obertrias) unter besonderer Berücksichtigung der Ostrakoden, 1: Ostrakoden der Zlambach-Schichten, besonders Bairdiidae. *Senckenbergiana Lethaea* 52: 129–283.
- Bradfield HH. 1935. Pennsylvanian ostracods of the Ardmore Basin, Oklahoma. *Bulletin of American Paleontology* 22: 1–145.
- Brady GS. 1880 Report on the Ostracoda dredged by H.M.S. Challenger during the Years 1873–1876. Report on the Scientific

- Results of the Voyage of H.M.S. Challenger during the years 1873–76. *Zoology* 1(Part 3): 1–184.
- Bunza G, Kozur H. 1971. Beiträge zur Ostracodenfauna der tethyalen Trias. *Geologisch Paläontologische Mitteilungen Innsbruck* 1: 1–76.
- Cafiero B, De Capoa Bonardi P. 1982. Biostratigrafia del Trias pelagico della Sicilia. *Bollettino della Società Paleontologica Italiana* 21(1): 35–71.
- Calcara P. 1840. Monografie dei generi Clausilia e Bulimo, coll'aggiunta di alcune nuove specie di conchiglie siciliane esistenti nella collezione della Sig. Teresa Gargotta in Salinas, 54 p.
- Calcara P. 1845. Cenno sui molluschi viventi e fossili della Sicilia da servire di supplemento ed insieme di critiche osservazioni all'opera di R.A. Philippi. Reale stamperia e Libreria Palermo, 65 p.
- Carrillat A. 2001. Palaeoenvironmental reconstruction of the Mufara Formation (Upper Triassic, Sicily): biostratigraphy, organic facies, sedimentological and geochemical approach. Thèse Université de Genève, Terre et Environnement 27: 269 p.
- Carrillat A, Martini R. 2009. Palaeoenvironmental reconstruction of the Mufara Formation (Upper Triassic, Sicily): High resolution sedimentology, biostratigraphy and sea-level changes. *Palaeogeography, Palaeoclimatology, Palaeoecology* 283: 60–76.
- Catalano R, D'Argenio B. 1982. Schema geologico della Sicilia occidentale. In: Catalano R, D'Argenio B, eds. *Guida alla geologia della Sicilia occidentale. Guide geologiche regionali*. Memorie della Società Geologica Italiana suppl. A2: 9–41.
- Chen DQ, Shi CG. 1982. Latest Permian ostracoda from Nantong, Jiangsu and from Miannyang, Hubei. Bulletin of the Nanjing Institute of Geology and Palaeontology. *Academia Sinica* 4: 105–152 (in Chinese).
- Crasquin S, Horne DJ. 2018. The palaeopsychrosphere in the Devonian. *Lethaia* 51: 547–563.
- Crasquin S, Sciuto F, Reitano A. 2018. Late Carnian (Tuvalian, Tropites dilleri zone) ostracods (Crustacea) from the Mufara Formation (Monte Scalpello, Central-Eastern Sicily, Italy). *Annales de Paléontologie* 104: 129–142.
- Crasquin-Soleau S, Grădinaru E. 1996. Early Anisian ostracode fauna from the Tulcea unit (Cimmerian North Dobrogean orogeny, Romania). *Annales de Paléontologie* 82: 59–116.
- Dana J. 1853. Tribe III: Cyproidea, Ostracoda. In: *Crustacea of United States Exploring Expedition during the Years 1838, 1839, 1840, 1841, 1842, under the Command of Charles Wilkes, U.S.N., with Atlas of 96 plates*, 13(2), 1277–1304. Philadelphia: C. Sherman.
- Delo DM. 1930. Some Upper Carboniferous ostracods from the Shale Basin of Western Texas. *Journal of Paleontology* 4: 152–178.
- Di Paolo L, Aldega L, Corrado S, Mastalerz M. 2012. Maximum burial and unroofing of Mt. Judica recess area in Sicily: implication for the Apenninic–Maghrebic wedge dynamics. *Tectonophysics* 530–531: 193–207.
- Di Stefano P, Favara R, Luzio D, Renda P, Cacciatori MS, Calò M, *et al.* 2015. A regional-scale discontinuity in western Sicily revealed by a multidisciplinary approach: A new piece for understanding the geodynamic puzzle of the southern Mediterranean. *Tectonics* 34: 1–19.
- Drexler E. 1958. Foraminiferen und Ostracoden aus dem Lias von Siebeldingen/Pfalz. *Geologisches Jahrbuch* 75: 475–554.
- Egorov VG. 1950. Ostrakody franskogo iarusu Russkoi platformy, 1. Kloedenellidae: Moscow-Leningrad Vses. Neft. Nauchno-Issled. Geol. Razved. Inst., Moscow, Filial, 140 p.
- Forel M-B, Crasquin S. submitted. Bounded by crises: an updated view of the evolution of marine ostracods during the Triassic. *Marine Micropalaeontology*.
- Forel M-B, Grădinaru E. 2018. First report of Ostracods (Crustacea) associated with Bithynian (Anisian, Middle Triassic) Tubiphytes-microbial reef in the North Dobrogean Orogen (Romania). *Papers in Palaeontology* 4(2): 211–244.
- Forel M-B, Tekin UK, Okuyucu C, Bedi Y, Tuncer A, Crasquin S. 2017. Discovery of a long-term refuge for ostracods (Crustacea) after the end Permian extinction: a unique Carnian (Late Triassic) fauna from the Mersin Melange, southern Turkey. *Journal of Systematic Palaeontology* 17(1): 9–58.
- Forel M-B, Kolar-Jurkovsek T, Jurkovsek B. 2019a. Ostracods from the “Raibl Beds” (Carnian, Late Triassic) of Belca section in Karavanke Mountains, Northwestern Slovenia. *Geodiversitas*, in press.
- Forel M-B, Thuy B, Wisshak M. 2019b. Digging into the ancestral stocks of Jurassic lineages: ostracods (Crustacea) from Carnian (Late Triassic) sponge mounds from the Maantang Formation (South China). *Bulletin de la Société Géologique de France* 190(9): 29.
- Gemmellaro GG. 1904. I Cefalopodi del Trias Superiore della regione occidentale della Sicilia. *Giornale di Scienze Naturali ed Economiche di Palermo* 24: 319.
- Gerry E, Honigstein A, Rosenfeld A, Hirsch F, Eshet Y. 1990. The Carnian salinity crisis: ostracods and palynomorphs as indicators of palaeoenvironment. In: Whatley R, Maybury C, eds. *Ostracoda and global events*. London: Chapman & Hall, pp. 87–99.
- Gramm MN. 1970. Otpchatki adduktora Triasovyykh cytherellid (Ostracoda) Primor'ya i nekotorye teorii filembriogeneza. *Paleontologicheskii Zhurnal* 1: 88–103.
- Grasso M. 2001. The apenninic-Maghrebic orogen in southern Italy, Sicily and adjacent areas. In: Vai GB, Martini IP, eds. *Anatomy of an Orogen: The Apennines and Adjacent Mediterranean Basins*. Kluwer Academic Publication GB, pp. 255–286.
- Gründel J. 1962. Zur Taxonomie der Ostracoden der Gattendorfia Stufe Thüringens. *Freiderger Forschungshilfe* C 151: 51–105.
- Gründel J. 1964. Zur Gattung *Healdia* (Ostracoda) und zu einigen verwandten Formen aus dem unter Jura. *Geologie* 13: 1–469.
- Gründel J. 1967. Zur Grossgliederung der Ordnung Podocopida G.W. Müller, 1894. *Neues Jahrbuch für Geologie and Palaontologie, Monatshefte* 6: 321–332.
- Guernet C, Lethiers F. 1989. Ostracodes et recherche des milieux anciens : possibilités et limites. *Bulletin de la Société Géologique de France* 8(5): 577–588.
- Gümbel CW. 1869. Über Foraminiferen, Ostracoden und mikroskopische Tier-Ueberreste in den St. Cassianer und Raibler Schichten. *Jahrbuch Geologischen Reichsanstalt* 19: 175–186.
- Harlton BH. 1933. Micropaleontology of the Pennsylvanian Johns Valley Shale of Ouachita Mountains, Oklahoma and its relationships to the Mississippian Caney Shale. *Journal of Paleontology* 7: 3–29.
- Hirsch F, Gerry E. 1974. Conodont and Ostracode-biostratigraphy of the Triassic in Israel. *Schriftenreihe der Erdwissenschaftlichen Kommissionen, Österreichische Akademie der Wissenschaften* 2: 107–112.
- Horne DJ, Cohen A, Martens K. 2002. Taxonomy, morphology and biology of Quaternary and living Ostracoda. In: Holmes JA, Chiva AR, eds. *The Ostracoda: Applications in Quaternary Research*. American Geophysical Union, Geophysical Monograph 131: 5–36.
- Kaufmann A. 1900. Über zwei neue Candona-Arten aus der Schweiz. *Zoologischer Anzeiger* 23: 108–110.
- Keim L, Brandner R, Krystyn L, Mette W. 2001. Termination of carbonate slope progradation: An example from the Carnian of the Dolomites, Northern Italy. *Sedimentary Geology* 143: 303–323.
- Klie W. 1938. Ostracoda, Muschelkrebse. In: *Dahl, Die Tierwelt Deutschlands*. Jena 34(3): 1–230.
- Kollmann K. 1960. Ostracoden aus der alpinen Trias Österreichs. I Parabairdia n. g. und Ptychobairdia n. g. (Bairdiidae). *Jahrbuch der Geologischen Bundesanstalt* 5: 79–105.

- Kollmann K. 1963. Ostracoden aus der alpinen Trias. II. – Weitere Bairdiidae. *Jahrbuch der Geologischen Bundesanstalt Wien* 106: 121–203.
- Kornicker LS, Sohn IG, 1976. Phylogeny, ontogeny, and morphology of living and fossil Thaumatozoidacea (Myodocopa: Ostracoda). *Smithsonian Contribution to Zoology* 219: 1–124
- Kozur H. 1970. Neue Ostracoden-Arten aus dem Obersten Anis des Bakonyhochlandes (Ungarn). *Berichte des naturwissenschaftlich-medizinischen Vereins in Innsbruck* 58: 384–428.
- Kozur H. 1971a. Die Bairdiacea des Trias. Teil 1: Skulpturierte Bairdiidae aus Mitteltriassischen Flachwasse Ablagerungen. *Geologisch Paläontologische Mitteilungen Innsbruck* 1: 1–27.
- Kozur H. 1971b. Die Bairdiacea der Trias. Teil 2: Skulpturierte Bairdiidae aus Mitteltriassischen tiefschelfablagerungen. *Geologisch Paläontologische Mitteilungen Innsbruck* 1(5): 1–21.
- Kozur H. 1971c. Die Bairdiacea der Trias. Teil 3: Einige neue Arten triassischer Bairdiacea und Bemerkungen zur Herkunft der Macrocyprididae (Cypridacea). *Geologisch Paläontologische Mitteilungen Innsbruck* 1: 1–18.
- Kozur H. 1973. Beiträge zur ostracodenfauna der Trias. *Geologisch Paläontologische Mitteilungen Innsbruck* 3(5): 1–41.
- Kozur H, Nicklas L. 1970. Ostrakoden aus dem Plattenkalk-Niveau des Haupt Dolomites (Rheintal). In: Mostler H, ed. *Beiträge zur Mikrofazies und Stratigraphie von Tirol und Vorarlberg*. Festband des Geologischen Institutes, 300-Jahr-Feier Universität Innsbruck, pp. 309–320.
- Kristan-Tollmann E. 1970. Einige neue Bairdien (Ostracoda) aus der alpinen Trias. *Neues Jahrbuch für Geologie und Paläontologie* 135: 268–310.
- Kristan-Tollmann E. 1971. Weitere Beobachtungen an skulptierten Bairdiidae (Ostracoda) der alpinen Trias. *Neues Jahrbuch für Geologie und Paläontologie* 139: 57–87.
- Kristan-Tollmann E. 1972. Die obertriadischen Arten der Ostracodengattung *Kerocythere* Kozur and Nicklas 1970, und ihr stratigraphischer Wert. *Erdoel-Erdgas-Zeitschrift* 88: 43–49.
- Kristan-Tollmann E. 1973. Zur phylogenetischen und stratigraphischen Stellung der triadischen Healdiiden (Ostracoda) II. *Erdoel-Erdgas-Zeitschrift* 89: 150–155.
- Kristan-Tollmann E. 1977a. On the development of the muscle-scar patterns in Triassic Ostracoda. In: Löffler H, Danielopol D, eds. In: *Proceedings of the Sixth International Symposium on Ostracoda* (Saalfelden, Salzburg), pp. 133–143.
- Kristan-Tollmann E. 1977b. Zur Evolution des Schließmuskelfeldes bei Healdiidae und Cytherellidae (Ostracoda). *Neues Jahrbuch für Geologie und Paläontologie* 10: 621–639.
- Kristan-Tollmann E. 1978. Bairdiidae (Ostracoda) aus den obertriadischen Cassianer Schichten der Ruones-Wiesen bei Corvara in Südtirol. *Schriftenreihe der Erdwissenschaftlichen Kommissionen. Österreichische Akademie der Wissenschaften* 4: 77–104.
- Kristan-Tollmann E. 1983. Ostracoden aus dem Oberanien von Leidapo bei Guiyang in Süd China. *Neue Beiträge zur Biostratigraphie der Tethys-Trias* 121–176.
- Kristan-Tollmann E. 1991. Ostracods from the Middle Triassic Sina Formation (Aghdarband Group) in NE-Iran. *Abhandlungen der Geologischen Bundesanstalt* 38: 195–200.
- Kristan-Tollmann E, Hamedani A. 1973. Eine spezifische Mikrofaunen-Vergesellschaftung aus den Opponitzer Schichten der Oberkarn der niederösterreichischen Kalkvoralpen. *Neues Jahrbuch für Geologie und Paläontologie* 143(2): 193–222.
- Kristan-Tollmann E, Tollmann A, Hamedani A. 1979. Beiträge zur Kenntnis der Trias von Persien. I. Revision der Triasgliederung, Rhäfazies im Raum von Isfahan und Kössener Faziesanschlag bei Waliabad SE Abadeh. *Mitteilungen der Gesellschaft der Geologie und Bergbaustudenten in Österreich* 70: 119–186.
- Latreille PA. 1806. Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata. Parisiis et Argentorati, 303 p.
- Lentini F. 1974. Caratteri stratigrafici e strutturali della zona di Monte Judica (Sicilia Orientale). *Rivista Mineraria Siciliana* 145-147: 71–96.
- Lentini F, Grasso M, Carbone S. 1987. *Introduzione alla geologia della Sicilia e guida all'escursione. Convegno della Società Geologica Italiana su sistemi avanssosa-avanpaese lungo la Catena Appenninico-Maghrebide, 22–25 aprile*, Naxos-Pergusa, Catania, 60 p.
- Lethiers F, Raymond D. 1991. Les crises du Dévonien supérieur par l'étude des faunes d'ostracodes dans leur cadre paléogéographique. *Palaeogeography, Palaeoclimatology, Palaeoecology* 88: 133–146.
- Lieberman HM. 1979. Die Bivalven- und Ostracodenfauna von Raibl und ihr stratigraphischer Wert. *Verhandlungen der Geologischen Bundesanstalt* 2: 85–131.
- Lord AR. 1982. Metacopine ostracods in the Lower Jurassic. In: Bannar ET, Lord AR, eds. *Aspects of Micropalaeontology*. London: Allen & Unwin, pp. 262–277.
- Lucas SG. 2010. The Triassic timescale: an introduction. In: Lucas SG, ed. *The Triassic Timescale*, 334. Geological Society, London, Special Publications, pp. 1–16.
- Maddocks RF. 1969. Revision of Recent Bairdiidae. *Bulletin of the United States National Museum* 295: 1–126.
- Mandelstam NI, Schneider GF, Kuznetsova ZV, Kats FI. 1957. [New genera of Ostracoda in the families Cypridae and Cytheridae]. All-Union Palaeontological Society, Annual. *Palentologičeskogo Obščestva Ezhegodnik* 16: 182–183 [in Russian].
- Maugeri Patané G. 1934. Sopra alcuni coralli del Trias superiore di M. Scalpello (Enna). *Relationes Consessuum Societatis Gioienae Catinensis Naturalium Scientiarum* 67: 9.
- McCoy F. 1844. A synopsis of the characters of the Carboniferous Limestone fossils of Ireland. Dublin: Dublin University Press, 207 p.
- Méhes G. 1911. Über Trias-Ostrakoden aus dem Bakony. Magyar Földrajzi Társaság, Budapest, Palaton-bizottság, pp. 1–38.
- Mette W, Mohtat-Aghai P. 1999. Ostracods and foraminifera from Upper Triassic intrashelf basin deposits in the Northern Calcareous Alps. *Geologisch Paläontologische Mitteilungen Innsbruck* (unpublished manuscript).
- Mette W, Honigstein A, Crasquin S. 2014. Deep-water ostracods from the Middle Anisian (Reifling Formation) of the Northern Calcareous Alps (Austria). *Journal of Micropalaeontology* 34: 71–91.
- Michelsen O. 1975. Lower Jurassic biostratigraphy and ostracods of the Danish Embayment. *Danmarks Geologiske Undersøgelse*, Raekke 2(104): 1–287.
- Monostori M. 1994. Ostracod evidence of the Carnian salinity crisis in the Balaton Highland, Hungary. *Neues Jahrbuch für Geologie und Paläontologie* 193: 311–331.
- Monostori M. 1995. Environmental significance of the Anisian Ostracoda fauna from the Forras Hill near Felsoors (Balaton Highland, Transdanubia, Hungary). *Acta Geologica Hungarica* 39: 37–56.
- Monostori M, Tóth E. 2013. Ladinian (Middle Triassic) silicified ostracod faunas from the Balaton Highland (Hungary). *Rivista Italiana di Paleontologia e Stratigrafia* 119: 303–323.
- Monostori M, Tóth E. 2014. Additional Middle to Upper Triassic ostracod faunas from the boreholes of Transdanubian Central Range (Hungary). *Hantkeniana* 9: 21–43.

- Montanari L. 1987. Lineamenti stratigrafico-paleogeografici della Sicilia durante il Ciclo Alpino. *Memorie della Società Geologica Italiana* 38: 361–406.
- Moore RC. 1961. Treatise of Invertebrate Paleontology. Part Q. Arthropoda 3, Crustacea, Ostracoda. Geological Society of America and University of Kansas Press, 442 p.
- Müller GW. 1894. Die Ostracoden des Golfes von Neapel und der angrenzenden Meere Abschnilte. *Fauna und Flora des golfes von Neapel* 21: 1–404.
- Müller GW. 1906. Die Ostracoden der Siboga-Expedition. In: Max W, ed. *Siboga-Expedition, Uitkomsten op Zoologisch, Botanisch, Oceanographisch en H.M. Siboga. Monographien* 30, EJ Brill, Leiden, 40 p.
- Nelli B. 1899a. Il Raibliano del Monte Judica nella provincia di Catania. *Atti della Reale Accademia Nazionale dei Lincei, Rendiconti della Classe di Scienze fisiche, matematiche e naturali* s. 5, 8(1): 91–92.
- Nelli B. 1899b. Il Raibli nei dintorni di M. Judica. *Bollettino della Società Geologica Italiana* 18(2): 211–223.
- Oertli HJ. 1971. The aspect of ostracode faunas – A possible new tool in petroleum sedimentology. *Bulletin du Centre de Recherches de Pau* 5(suppl.): 137–151.
- Özdkimen H. 2010. Substitute names for three genera of Ostracoda (Crustacea). *Munis Entomology and Zoology* 5: 315–316.
- Pokorny V. 1964. Some paleoecological problems in marine ostracode faunas, demonstrated on the Upper Cretaceous ostracodes of Bohemia, Czechoslovakia. *Pubblicazioni della Stazione Zoologica di Napoli* 33(Suppl.): 462–479.
- Reuss AE. 1869. Zur fossilen Fauna der Oligocän-schichten von Gaas. *Sitzungsberichte der österreichischen Akademie der Wissenschaften, mathematisch-naturwissenschaftliche Klasse* Abt.1. 59(3): 446–488.
- Salaj J, Jendrejakova O. 1984. Ecology and facial relation of some groups of Triassic foraminifers and ostracods of stratigraphic importance. *Geologický Zborník-Geologica Carpathica* 35(2): 231–240.
- Sars GO. 1866. Oversight af Norges marine Ostracoder. *Forhandlinger i Videnskabs-Selskabet I Christiania* 1865, pp. 1–130.
- Scalia S. 1907a. I fossili del Trias superiore dell'Acquanova e di Paraspora. *Bollettino dell'Accademia Gioenia di Scienze Naturali in Catania* s. 2(1-2): 10–12.
- Scalia S. 1907b. La fauna del Trias superiore del gruppo montuoso di Judica. *Bollettino dell'Accademia Gioenia di Scienze Naturali in Catania* s. 2(3-4): 17–25.
- Scalia S. 1909. Il gruppo di M. Judica. *Bollettino della Società Geologica Italiana* 28: 269–340.
- Scalia S. 1910–1914. La fauna del Trias superiore del gruppo del M. Judica: Parti 1–3. *Atti dell'Accademia Gioenia di Scienze Naturali in Catania* s. 5 (3-5-7): 51+58+25.
- Schmidt di Friedberg P, Trovò A. 1962. Contribution à l'étude structurale du groupe du Monte Judica (Sicile orientale). *Bulletin de la Société Géologique de France* 4(5): 754–759.
- Sebe O, Crasquin S, Grădinaru E. 2013. Early and Middle Anisian (Triassic) deep-water ostracods (Crustacea) from North Dobrogea (Romania). *Revue de Paléobiologie* 32: 509–529.
- Sohn IG. 1968. Triassic Ostracodes from Makhtesh Ramon, Israel. *Bulletin of the Geological Survey of Israel* 44: 1–71.
- Speranza F, Minelli L. 2014. Ultrathick Triassic dolomites control the rupture behavior of the central Apennine seismicity: Evidence from magnetic modeling of the L'Aquila fault zone. *Journal of Geophysical Research: Solid Earth* 119: 1–15.
- Styk O. 1958. Triassic microfauna in the neighbourhood of Chrzanow and the northwestern part of the Mesozoic periphery of the Swiety Kryzysz Mts. *Bulletin Poland Instytut Geologiczny* 121, *Micropalaeontological researches in Poland* 3: 163–176.
- Sylvester-Bradley PC. 1961. Suborder Metacopina Sylvester-Bradley, n. suborder. Q358–359. In: Moore RC, ed. *Treatise of invertebrate paleontology. Part Q, Arthropoda 3, Crustacea, Ostracoda*. Lawrence, KS: The Geological Society of America, Boulder, CO and University of Kansas Press, 442 p.
- Széles M. 1965. Ostracodák a Bakonyi Nosztori-Volgy felsokarni rétegeibol. *Földtani közlöny* 45: 412–417.
- Tollmann A. 1976. Analyse des klasschen nordalpinen Mesozoikums. *Monographie des Nördlichen Kalkalpen* 2: 1–580.
- Triebel E. 1941. Zur Morphologie und Ökologie der fossilen Ostracoden. Mit Beschreibung einiger neuer Gattungen und Arten. *Senckenbergiana* 23: 294–400.
- Ulrichs M. 1970. Variability of some Ostracods from the Cassian beds (Alpine Triassic) depending on the ecology. *Bulletin du Centre Recherches de Pau – SNPA* 5: 695–715.
- Zorn I. 2010. Ostracodal Type Specimens Stored in the Paleontological Collection of the Geological Survey of Austria. *Jahrbuch der Geologischen Bundesanstalt* 150: 263–299.

**Cite this article as:** Crasquin S, Sciuto F, Reitano A, Coco RM. 2020. Late Triassic (Tuvallian – Carnian, *Tropites subbullatus*/*Anatropites spinosus* zones) ostracods from Monte Gambanera (Castel di Iudica, Central-Eastern Sicily, Italy), *BSGF - Earth Sciences Bulletin* 191: 36.