

# The transition from Pyrenean shortening to Gulf of Lion rifting in Languedoc (South France) – A tectonic-sedimentation analysis

Michel Séranne<sup>1,\*</sup>, Renaud Couëffé<sup>2</sup>, Eglantine Husson<sup>2</sup>, Céline Baral<sup>1</sup> and Justine Villard<sup>1</sup>

<sup>1</sup> Géosciences Montpellier, Université de Montpellier – CNRS, Montpellier, France

<sup>2</sup> BRGM, GeoResources Division, Orléans, France

Received: 29 July 2020 / Accepted: 9 April 2021 / Publishing online: 4 May 2021

**Abstract** – The Pyrenean orogen extended eastward, across the present-day Gulf of Lion margin. The late or post-orogenic dismantling of this orogen segment, contemporaneous with ongoing shortening in the Spanish Pyrénées, is still debated. Understanding the transition between the two geodynamic events requires to document the precise timing of the succession of the tectonic processes involved. We investigate the superposition of rifting structures over Pyrenean thrusts and folds in the onshore Languedoc. Compilation and reassessment of the regional chronostratigraphy, in the light of recent biostratigraphic dating and new mapping of Paleogene basins, lead to date the transition to the Priabonian. Tectonic-sedimentation relationship in the Eocene to Oligocene depocentres are analysed in surface exposures as well as in seismic reflection surveys. Bed-to bed mapping allowed us to: i) characterise an intermediate sequence of Priabonian age, bounded at the base and the top by unconformities; ii) evidence syn-depositional deformation within the Priabonian; iii) define the axes of Priabonian deformation. Interpretation of seismic reflection profiles, across the onshore basins covered by syn- and post-rift sequences, reveals the existence of an intermediate sequence displaying similar features, and that is correlated to the Priabonian. Syn-depositional deformation of some Priabonian basins correspond to extensional structure, whereas neighbouring, contemporaneous basins, reveal compressional deformation. The distribution of such apparently conflicting observations across the studied area provides evidence for left-lateral strike-slip deformation between two major regional faults (Cévennes and Nîmes faults). Left-lateral strike-slip along NE-trending faults accommodates E-W extension of the West European Rift (ECRIS) and part of the ongoing N-S shortening in the Central and Western Pyrénées. Priabonian clastic sedimentation and deformation in Languedoc witness the initial stages of the dismantling of the Languedoc-Provence Pyrénées, prior to Oligocene-Aquitainian back-arc rifting.

**Keywords:** Pyrénées / Gulf of Lion / strike-slip basins / syn-depositional deformation / Priabonian

**Résumé** – Transition entre la compression Pyrénéenne et l'extension du Golfe du Lion, dans le Languedoc (Sud de la France) par analyse des relations tectonique-sédimentation. L'orogène pyrénéen s'étendait vers l'est à l'emplacement actuel du Golfe du Lion. Contemporain de la poursuite de la compression dans les Pyrénées franco-espagnoles, le démantèlement tardi- ou post-orogénique de ce segment de l'orogène est encore mal compris. La compréhension de la transition entre l'orogénèse Pyrénéo-Provençale et le rifting du Golfe du Lion nécessite de préciser le calendrier des événements ainsi que la succession des différents processus tectoniques mis en jeu. Nous analysons la superposition des structures de rifting sur les structures compressives Pyrénéennes exposées à terre, dans le Languedoc. Tout d'abord, nous compilons et révisons la chronostratigraphie régionale, à la lumière de datations postérieures à la publication des cartes au 1/50 000 du BRGM, et de la cartographie de plusieurs bassins paléogènes. Nous datons cette transition du Priabonien. Nous analysons les relations tectonique-sédimentation des dépocentres Eocène à Oligocène sur le terrain et en sismique réflexion. La cartographie banc-par-banc permet de : i) caractériser une séquence intermédiaire d'âge Priabonien encadrée par deux discordances, ii) mettre en évidence une déformation syn-dépôt dans le remplissage Priabonien, iii) définir les axes de la déformation priabonienne. L'interprétation de profils de sismique réflexion à travers les bassins paléogènes à terre, recouverts par les séries syn- et post-rift, révèle l'existence d'une séquence intermédiaire présentant des caractéristiques similaires et que nous corrélons avec le Priabonien. La déformation syn-dépôt de

\*Corresponding author: [michel.seranne@umontpellier.fr](mailto:michel.seranne@umontpellier.fr)

certaines bassins est extensive, alors que dans des bassins voisins contemporains, elle est compressive. La distribution régionale de ces observations apparemment contradictoires est interprétée comme liée à un couloir de déformation en décrochement senestre entre les failles majeures des Cévennes et de Nîmes, d'orientation NE-SW. Ce décrochement accommode l'extension EW des rifts ouest-européens et contribue au raccourcissement N-S des Pyrénées franco-espagnoles, toujours actif au Priabonien. La sédimentation syn-tectonique clastique du Priabonien du Languedoc enregistre les premiers stades du démantèlement de la chaîne pyrénéo-provençale, plusieurs millions d'années avant l'initiation de l'extension d'arrière-arc.

**Mots clés** : Pyrénées / Golfe du Lion / bassins sur décrochement / déformation syn-sédimentaire / Priabonien

## 1 Introduction

The Pyrenean orogen results from the inversion of Mesozoic extensional basins, mostly formed during Mid-Cretaceous rifting and mantle denudation (Vergés and Garcia-Senz, 2001; Lagabrielle and Bodinier, 2008; Lagabrielle *et al.*, 2010; Tugend *et al.*, 2014; Chelalou *et al.*, 2016; Cochelin *et al.*, 2018; Ternois *et al.*, 2019). The orogenic shortening was accommodated in the upper crust by reactivation of the inherited Cretaceous extensional faults, as thrust faults (Debroas, 1990), while the middle crust and exhumed mantle were subducted beneath the European crust (Teixell *et al.*, 2016, 2018). The orogen extended eastwards (Ford and Vergés, 2020), in Corbières (Viallard, 1987; Lamotte *et al.*, 2002), Languedoc (Arthaud and Laurent, 1995) and southern Provence (Lacombe and Jolivet, 2005; Bestani *et al.*, 2015), across the present-day Gulf of Lion (Arthaud and Séguret, 1981; Gorini *et al.*, 1994). It was later dismantled in Oligocene time to give way to the Gulf of Lion rifted margin (Arthaud *et al.*, 1981; Arthaud and Séguret, 1981; Gorini *et al.*, 1994; Séranne *et al.*, 1995; Mauffret and Gorini, 1996; Mascle and Vially, 1999; Séranne, 1999; Guennoc *et al.*, 2000; Lacombe and Jolivet, 2005). In the onshore Corbières area, upper crust extension is characterised by “negative inversion” of the Pyrenean thrusts, associated with deposition of Oligocene-Aquitainian syn-rift sediments in the hanging-wall (Gorini *et al.*, 1991). The crustal roots of the Pyrénées disappear eastwards (Chevrot *et al.*, 2018) and the orogen brutally gives way to the Gulf of Lion rift and margin, within only tens of kilometres (Mauffret *et al.*, 2001; Jolivet *et al.*, 2020). Although documented both onshore and offshore, tectonics and kinematics of the superimposition of Pyrenean structures by later rifting (Fig. 1) remains incompletely understood. Transition of the compression to extension could be driven by collapse of the elevated mountain range (Séranne *et al.*, 1995), it could be related to the southern propagation of the European Cenozoic Rift System (ECRS) (Dèzes *et al.*, 2004), or to the onset of back-arc extension due to the NW-dipping subduction roll-back of the old Tethyan lithosphere (Séranne, 1999; Jolivet *et al.*, 2020). Better constraints on tectonics and precise timing of the transition from the Pyrenean orogeny to the Gulf of Lion rifting are needed in order to better understand the origin and driving mechanism of the transition between the two major geodynamics events. Such data are also needed to constrain paleotectonic restorations (*e.g.* Bestani *et al.*, 2015; Christophoul *et al.*, 2016), and kinematic reconstructions of the Mediterranean (*e.g.* Romagny *et al.*, 2020).

Since most of the collapsed orogen is presently buried beneath the thick Neogene sequence of the Gulf of Lion margin, the onshore Languedoc provides the only geological

outcrops that are relevant to address this question. There, the rare available stratigraphic data suggested that the Pyrenean orogeny lasted until “Early Oligocene”, as reported on the Saint-Martin-de-Londres 1/50 000-scale geological map (Philip *et al.*, 1978), while rifting took place in the late Rupelian according to the Montpellier map (Andrieux *et al.*, 1971). Consequences of such a short delay implies: i) that Pyrenean compression was polyphase, including a last compressive phase, following a Late Eocene period of tectonic quiescence, and ii) rifting interrupted folding and thrusting, with no transition.


In this paper, we examine five Paleogene basins located in Languedoc, which developed at the front of the Eocene external thrusts of the Pyrenean belt and onshore of the Gulf of Lion passive margin. The main objective of this contribution is to document, thanks to the revision of available biostratigraphic data and to detailed sedimentological, tectonic and seismic reflection analyses, the key period of the transition between the last stages of Pyrenean orogenic (mountain building) events and the earliest stages of rifting associated with the opening of the Gulf of Lion. We aim at constraining the tectonic setting associated with the deposition of Priabonian continental units by analysing the tectonic-sedimentation relationships in 5 distinctive basins (Fig. 2). Analyses are based on field observations (Guzargues, Les-Matelles, Saint-Martin-de-Londres basins) and subsurface data (Hérault and Saint-Chaptes-Gardon basins). This study also relies on: i) new original investigation in the Guzargues basin, ii) reappraisal of published works in Les-Matelles and Saint-Martin-de-Londres basins (the latter involving additional mapping), and iii) the interpretation of ancient (1983–1992) seismic profiles tied to boreholes data in the Hérault and Saint-Chaptes-Gardon basins.

## 2 Regional geological setting

### 2.1 Location of the studied area within the Pyrenean orogen

Languedoc, in the south of France, is located between the Variscan basement of the Massif Central and the shores of the Mediterranean Sea; it links the Pyrénées (in the SW) to the Alps (to the NE), across the Rhône Valley (Fig. 1). Together with the Corbières, Languedoc forms a segment of the north foreland of the Pyreneo-Provençal fold and thrust belt (Arthaud and Séguret, 1981; Bestani *et al.*, 2015).

The Paleozoic (Variscan) basement is exposed in the Massif Central to the NW, in the Maure and Esterel massifs to the east. To the south, the basement has been reactivated during the Late Cretaceous to Eocene Pyrenean orogeny. It is exposed



**Fig. 1.** Structural map southern France showing the relation between the Pyrénées and Gulf of Lion margin; modified from (Séranne *et al.*, 1995). Na: Narbonne; Pe: Perpignan; Mo: Montpellier; Ma: Marseille. Directions of Pyrenean shortening from Gaviglio and Gonzales (1987), Lacombe *et al.* (1992), Arthaud and Laurent (1995) and Lamotte *et al.* (2002) and Oligocene extension from Arthaud *et al.* (1977), Benedicto (1996) and Hippolyte *et al.* (1993).


in the Pyrénées and buried beneath the thick Neogene sequences of the Gulf of Lion margin (Arthaud and Séguret, 1981; Gorini *et al.*, 1994; Guennoc *et al.*, 2000) (Fig. 1). Onshore, the Variscan basement is unconformably covered by Mesozoic sequence, including at the base variable amounts of Triassic detritals and evaporites (Debrand-Passard and Courbouleix, 1984), which have been used as decollement level during later tectonic phases. The Jurassic through to Neocomian marine marls and carbonates were deposited during the Tethyan rifting and subsequent thermal subsidence, controlled by NE-trending normal faults such as the Cévennes, Nîmes and Durance faults. Thickness of the Jurassic increases southeastwards from the Cévennes fault (less than 1.5 km), and maximum subsidence is recorded in the hanging-wall of the Nîmes Fault, where seismic profiles show up to 10 km of Trias to Neocomian sequences (Séguret *et al.*, 1997; Le Pichon *et al.*, 2010).

This Mesozoic sedimentary cover has been folded and thrust in the Pyrenean foreland, detached over Triassic decollement. Thick-skinned thrusting presumably occurred close to the present day shoreline (Arthaud and Séguret, 1981; Séranne *et al.*, 1995; Lacombe and Jolivet, 2005). The style of the thrust and fold belt is mostly controlled by the thickness of the detached Mesozoic sedimentary cover (Arthaud and Laurent, 1995): north-verging, EW-oriented thrusts dominate areas of thin cover ( $\leq 2.5$  to 3 km) whereas thicker cover

developed series of E-W folds. Combined thin- and thick-skinned shortening exceed 20 km in the Languedoc (Arthaud and Laurent, 1995) and 40 km in the Provence segment of the orogen (Bestani *et al.*, 2015, 2016). The Pyrénées-Provence fold and thrust belt is segmented by NE- to NNE-trending faults which separate areas of distinct structural styles and amounts of shortening. Left-lateral strike-slip along the Cévennes Fault accommodated shortening of its hanging-wall while the NW footwall remained mostly undeformed during the Pyrenean orogeny (Arthaud and Laurent, 1995). In Provence, the Durance Fault separates thin- and thick-skinned Pyrenean deformation (Bestani, 2015; Bestani *et al.*, 2016).

## 2.2 Location of the studied area within the Gulf of Lion rift and passive margin

The studied area is affected by NE-trending normal faults bounding syn-rift Oligocene basins (Fig. 2). At least, some of these faults – if not all – are reactivated previous Pyrenean left-lateral strike-slips faults (*i.e.* the Cévennes fault, Matelles Fault (Mattaier, 2002). Seismic reflection profiling has shown that, during Oligocene rifting, such faults detached in the Triassic decollement (Benedicto, 1996; Benedicto *et al.*, 1999; Sanchis and Séranne, 2000; Husson, 2013). The onshore syn-rift basins NW of the Nîmes Fault were formed



**Fig. 2.** Simplified geological map of Languedoc, in the northern foreland of the Provence-Pyrénées fold and thrust belt. See location in the regional structural framework of [Figure 1](#). Priabonian basins according to the present study, in some areas, stratigraphic attributions are different from the BRGM 1/50 000-scale geological maps (see text for explanation). Subsurface data used in the study is indicated fine black lines (seismic) and open circles (boreholes); seismic profiles shown in this contribution are in bolder black line. Location of detailed maps and sections is shown with double black line frames.

by thin-skinned extensional tectonics (Séranne, 1999). The basement-cover decollement ramps down in the Nîmes basement fault (Benedicto *et al.*, 1996) and into the basement ramp of the Montpellier thrust to the south and southeast of the study area.


### 3 Revised chronology of events

The chronology of the transition between Pyrenean orogeny and the Gulf of Lion rifting is still discussed. The stratigraphy of the 1/50 000-scale geological maps of the study area is confusing: all maps are consistent in identifying the Bartonian (e6) as a Pyrenean syn-tectonic sequence, but they

greatly diverge in dating the end of the orogeny and the onset of the rifting. The comparison of stratigraphy of the adjacent maps of the study area reveals contradicting ages for identical and spatially continuous formations (Fig. 3a).

Another inconsistency arises from the outstanding question: what was the geodynamic context during the time interval between the Pyrenean syn-orogenic deposits and the syn-rift formations? Adjacent maps carry contradicting interpretations: the “g1” is a continuity of the syn-orogenic Bartonian on the Saint-Martin-de-Londres map (Philip *et al.*, 1978), whereas the Montpellier map correlates the “g1” with the early stages of the Gulf of Lion rifting (Andrieux *et al.*, 1971).

The mammalian reference levels developed for the Paleogene continental record of Europe (Sigé and Legendre,



**Fig. 3.** a: Correlation chart of the Eocene to Oligocene interval, and inferred ages, according to the BRGM 1/50 000-scale geological maps of the study area. The inset in the lower part gives the position of the BRGM 1/50 000-scale geological maps. b: Revised chronostratigraphy of the basins north of Montpellier (covered by the geological maps of Montpellier, St Martin-de-Londres and part of Sommières and Anduze) and of the Saint-Chaptes-Gardon basins (maps of Alès and part of Anduze and Uzès). This is based on new mapping of the Paleogene basins (this study) and on the compilation of biostratigraphy data, including recent discoveries of mammal fossils (references in the figure and in the text). Biostratigraphy scale according to mammal reference levels (M.Z.). Black stars and black hexagons correspond to mammal and charophyte sites, respectively. Simplified lithology of the formations is indicated. Colour code as in the following maps and sections of the Paleogene basins analysed in this study. White areas correspond to non-depositional and erosional hiatuses; the Priabonian interval from Saint-Chaptes-Gardon-Alès basins is documented in [Lettéron \*et al.\* \(2018\)](#).

1997), allowed to improve the stratigraphy and to revise the tectonic agenda of the area, especially around the transition from orogeny to rifting. [Figure 3b](#) uses chronology of [Vandenbergh \*et al.\* \(2012\)](#) and mammal reference levels of [Aguilar \*et al.\* \(1997\)](#).

The age of the lacustrine limestone was defined as Lutetian or slightly older. In the Pic-Saint-Loup foreland, fauna allowed to place this formation between MP12 and MP13 reference levels of the mammal reference levels ([Crochet \*et al.\*, 1988](#)). In different localities of Les-Matelles basin, several lignite beds provided a diversified fauna, which correlates with the Lutetian (MP13) ([Crochet \*et al.\*, 1997](#)). A distinctive marly facies including 1 to 20 cm diameter oncoliths, occurs at the top. It has yielded fauna ascribed to the MP14 reference level ([Crochet \*et al.\*, 1997](#)) of latest Lutetian age. This facies passes southwards and upwards to the Bartonian syn-orogenic breccia, exposed along the Montpellier thrust ([Andrieux \*et al.\*, 1971](#)). In the Sommières and Saint-Chaptes basins the

reference sites of Robiac provided a late Bartonian age for the latest Pyrenean syn-tectonic formations ([Remy, 2015](#)).

They pass upward to continental marls and limestones dated to the Bartonian-Priabonian transition (MP17) ([Remy, 2015](#); [Rémy and Lesage, 2005](#)). In the Saint-Martin-de-Londres basin, similar lithologies correspond to time equivalent, as suggested by charophytes (Feist in ([Philip \*et al.\*, 1978](#))).

The overlying formation mapped as “g1” in Les-Matelles basin, displays polygenic conglomerates and sandstones within yellow marls and silts, including several lignite beds. They are interpreted as alluvial plain deposits ([Benedicto \*et al.\*, 1999](#); [Egerton, 1996](#)). The age of this formation was subjected to discussion due to the controversial location of the discovery by Gervais, in the late XIXth Century, of a *Lophiodon* skull (see review in ([Hartenberger \*et al.\*, 1969](#))). Reappraisal of the specimen, exhumed from Lyon university collections by J.A. Remy (*personal communication*), relates the specimen to




Fig. 4. Generalised lithostratigraphy of the Late Mesozoic and Paleogene in the Guzargues Basin and immediate surrounding.

MP18 (middle Priabonian). Another *Lophiodon* specimen has recently been discovered in the correlative “g1” formation in the Guzargues basin, and has been related to the MP19 reference level (R. Tabuce and F. Lihoreau, *personal communication*). The age of this conglomerate formation is therefore attributed to the Priabonian. As first suggested by [Andrieux \*et al.\* \(1971\)](#), it is correlated with the detrital continental formation “Grès de Cèlas” of upper Priabonian, in the Sommières and Saint-Chaptes-Gardon basins, northeast of the study area, which spans the MP18 to MP19 ([Rémy, 1985](#); [Remy and Fournier, 2003](#); [Rémy and Lesage, 2005](#)).

At Les-Matelles and Saint-Vincent-de-Barbeyrargues localities, yellow to orange marls interfingering with syn-rift breccia, contain charophytes ([Grambast, 1962](#)), teeth of small mammals ([Thaler, 1962](#)) and gastropods ([Rey, 1962](#)) that yielded mid- to late- “Stampan” age, which corresponds to the Late Rupelian. This age was later confirmed by the discovery of small mammals in Les-Matelles half-graben, that relate to MP25 mammal scale ([Crochet, 1984](#)). In addition, breccia pipes and dykes of intracontinental alkaline basalt, that intrude Lutetian and Priabonian formations, were dated 24 to 28 Ma ([Gastaud \*et al.\*, 1983](#)), and have been related to the Late Rupelian rifting event ([Séranne, 1999](#); [Dautria \*et al.\*, 2010](#)). In the northern basins, this unconformable formation has not yet provided mammal remains, but Charophytes from the alluvial environments of Saint-André-de-Cruzières in the north of Alès Basin yielded an Oligocene age ([Feist-Castel, 1971](#)).

Reappraisal of ancient specimens and new discoveries thus indicate that the sequence made of marls, sandstones beds and polygenic channelized conglomerates, which is exposed in separated basins across the study area, belongs to the same

interval, dated to the Priabonian. In particular, the fluvial conglomerate and yellow sandy marls (reported on the published maps as “g1”, [Fig. 3a](#)), correlate with, and are the proximal equivalent of the lacustrine delta sandstones “Grès de Cèlas” of the Alès Basin ([Lettéron \*et al.\*, 2018](#)). This formation is intermediate between the syn-orogenic Pyrenean (Bartonian) and the syn-rift (Late Rupelian) deposits. Furthermore, it is bounded by two marked angular unconformities, the upper one corresponding to a several-million-years hiatus.

In the following, we examine key field evidence in the north-Montpellier area (Guzargues, Les-Matelles and St-Martin-de-Londres basins, respectively) where the tectonic sedimentation relationships allow to characterise this intermediate formation. It can be interpreted as the record of the transition between the Pyrenean orogeny to the Gulf of Lion rifting.

## 4 Tectonic and sedimentation in the Guzargues Basin

### 4.1 Lithostratigraphy of Eocene deposits in the Guzargues area

The Guzargues basin ([Fig. 2](#)) has not been the topic of any specific study and remains poorly documented. The Paleogene sedimentary succession contains 4 main lithological units representative of the regional geodynamic history since the early stages of the Pyrenean compression to the main rifting phase ([Fig. 4](#)). When possible, all of them were precisely

mapped to analyse the geometrical relationships of Priabonian deposits with under- and overlying units.

- Paleocene to Lower Eocene facies are represented by fluvial marls with interbedded sandstones and palustrine limestones (“*Marnes infra-lutésiennes*”), only few tens of metres thick in the Guzargues area. They are usually correlated to alluvial fan syn-tectonic breccias, reworking the Jurassic succession at the front of the Montpellier thrust, and thus deposited during the early stages of the Pyrenean compression.
- Massive whitish Lutetian limestones, up to 100 m thick in the area, form an easily recognised, continuous unit displaying folds produced by late stages of the Pyrenean compression.
- Lutetian limestones are conformably capped by a singular unit, up to 30 m thick, made of lacustrine to shallow marine, marly facies containing oncoliths and calcarenitic layers, Bartonian in age.
- Topmost deposits of the Paleogene succession in the Guzargues area correspond to Priabonian fluvial deposits showing alternations of conglomerates, sandstones and marls, up to 300 m thick. Pebbles are supplied both from distant Pyrenean sources and from local Mesozoic carbonates. These deposits rest unconformably above the Pyrenean folds consisting of Lutetian to Bartonian units.

One kilometre to the south, the northern edge of the neighbouring Assas basin shows a sedimentary unit of strong geodynamic significance, which is missing in the Guzargues basin. Close to the Assas village, Oligocene breccias and marls, usually interpreted as syn-rift deposits, unconformably overlie Priabonian continental facies.

Thus, the Priabonian deposits of the Guzargues basin, bounded below and above by unequivocal syn-tectonic units, represent a key unit to study the transition between two main regional geodynamic events.

## 4.2 Sedimentological features of Priabonian deposits in the Guzargues basin

As observed in the 10 sections studied in the Guzargues basin, Priabonian deposits are basically made of eight lithofacies units distinguished by their lithology and sedimentary structures. Each lithofacies is described and interpreted in terms of depositional processes (Tab. 1), and most are illustrated in Figure 5.

Lithofacies units are often associated together, defining three main facies associations (FA1, FA2, and FA3), the distribution of which varies depending on the spatial location in the Guzargues basin and on the vertical position into the Priabonian stratigraphic succession.

**FA1** is represented by the alternations of (i) channel-shaped, poorly to weakly sorted conglomerates (Gu, Gt; Tab. 1), (ii) matrix-rich, stratified conglomerates (Gm; Tab. 1), and (iii) medium- to coarse-grained, cross-bedded sandstones (Sp; Tab. 1). A general trend in the vertical evolution of FA1 facies units is recognised through most of the studied sections. In the lower part of sections, predominant Gu/Gt conglomeratic channels, up to 3 m thick and 3 to 5 m wide, contain

heterometric, sub-rounded to well-rounded clasts, mainly composed of local Mesozoic (Upper Jurassic to Lower Cretaceous) carbonates. In some places, they also show abundant oncoliths (as fragments or complete specimens) probably originated from the underlying Bartonian formation. In the upper part of sections, conglomeratic bodies in Gm facies unit are thinner (< 1 m) and laterally more continuous (usually more than 10 m in width) than in the lower part. In addition to local Mesozoic carbonates, quartz and lydite, originated from Cévennes and/or Montagne Noire Paleozoic peripheral units, and ginger (Aptian?) sandstones represent a significant proportion of clasts in channel fillings.

**Gu** conglomerates can be interpreted as channel lag (Miall, 1977, 1978; Nemeč and Postma, 1993). The **Gm** and **Gt** facies units present strong similarities with gravel bars in braided streams (Miall, 1977; Nemeč and Postma, 1993; Mack and Leeder, 1999). **Sp** cross-bedded sandstones are typical of megaripples deposited by a unidirectional upper to lower flow regime (Simons *et al.*, 1965; Southard, 1991) produced by subaerial to subaqueous, stream and/or sheet flows (Mack and Leeder, 1999; Hampton and Horton, 2007). Inferred FA1 depositional environments are thought to belong to an alluvial fan setting, first supplied by a local catchment area (lower part of sections), and then by a regional catchment area (upper part of sections).

**FA2** corresponds to the alternation between (i) moderately sorted, medium- to coarse-grained sandstones with sub-rounded clast lenses (**Sst**; Tab. 1), and (ii) medium- to coarse-grained cross-bedded sandstones (**Sp**; Tab. 1). If local (Mesozoic but also Lutetian) calcareous elements still represent an important proportion of clasts (granules, gravels, pebbles), siliceous component (quartz and lydite gravels) is more abundant than in FA1. FA2 depositional processes are dominated by tractional transport under a unidirectional upper flow regime, with episodic sheet flows. Inferred depositional environments correspond to the proximal part of a subaqueous alluvial fan to fan delta, with sediment supplied from both local and regional catchment areas.

**FA3** is mainly represented by yellowish, greenish or reddish silty clays, with plastic texture at some places, in which are interbedded thin layers of well-sorted siltstones and fine-grained sandstones (**Fm**; Tab. 1). Carbonate pedogenic nodules locally form scarce, up to 1 m thick, distinctive layers (**P**; Tab. 1). Interbedded massive fine-grained sandstones (**Sr**; Tab. 1) usually show vertical root traces and more rarely asymmetrical ripple cross-laminations several centimetres thick at the top.

FA3 deposition must be the result of vertical settling from detrital suspensions in standing water (Horton and Schmitt, 1996; Miall, 1977). Fine-grained sandstones showing asymmetrical ripples and interbedded silty and sandy layers suggest episodic flow regime, related to sheet flood events. The occurrence of carbonate nodules records occasional pedogenic processes and temporary aerial exposure. Assuming these depositional processes, FA3 should have been deposited in the distal part of a subaqueous to subaerial alluvial fan, or in a floodplain laterally to major braided distributaries.

The 10 studied sections show vertical variations in the FA succession (basically FA1-FA2 or FA2-FA3 alternations) forming elementary sequences between several metres and tens of metres thick (Fig. 6). As indicated by the various inferred depositional environments for FA, these elementary

**Table 1.** Summary of the main facies features observed in the Priabonian deposits of the Guzargues basin, and interpretations in terms of depositional processes.

Facies code. Facies name	Lithology	Sedimentary structures	Depositional processes
<b>Gu.</b> Poorly sorted, massive, channel-shaped conglomerates (Fig. 7A)	Poorly sorted, massive, locally normally graded conglomerates. Highly heterometric, subrounded to well-rounded clasts (pebbles to small boulders), mainly composed of local Mesozoic (Upper Jurassic to Lower Cretaceous) carbonates.	Well-defined basal erosional surfaces with gutter/scour structures. Channels displaying few clast imbrications, 50 cm up to 3 m in depth, 3 to 5 m in width.	Channel-lag deposits under subaerial flashy braided-stream flows (Barrier <i>et al.</i> , 2010)
<b>Gt.</b> Weakly sorted, cross-bedded, stratified conglomerates with coarse-grained sand lenses (Fig. 7B)	Clast-supported, stratified conglomerates. Weakly sorted, heterometric, subrounded clasts (granules to small boulders of Mesozoic carbonates). Occurrence of poorly sorted lenses of coarse-grained sand matrix surrounding individual pebbles.	Fining upward infill. Trough cross-bedding.	Minor channel fills (Miall, 1978). Transverse- or diagonal-bar deposits under subaerial perennial braided-stream flows (Barrier <i>et al.</i> , 2010).
<b>Gm.</b> Matrix-rich, normally graded, stratified conglomerates (Fig. 7C)	Matrix-rich, normally graded, stratified conglomerates. Poorly sorted, subangular to subrounded clasts (granules to cobbles) made of local Mesozoic carbonates, common siliceous (quartz) clasts (originated from Cévennes and/or Montagne Noire Paleozoic peripheral units) and ginger (Aptian?) sandstones.	Some beds with horizontal alignment of clasts. Few clast imbrications into horizontal foreset strata. “Gradual” basal surfaces.	Longitudinal-bar deposits under subaerial competent and perennial braided-stream flows (Barrier <i>et al.</i> , 2010)
<b>Sst.</b> Moderately sorted, medium- to coarse-grained, cross-stratified sandstones with lenses of subrounded gravels and pebbles (Fig. 7C)	Moderately sorted, medium- to coarse-grained sandstones. Layers (and lenses) of well-sorted subrounded clasts (granules to pebbles) composed of Mesozoic carbonates, some Lutetian mudstones and common Bartonian oncoliths as fragments or complete specimens.	Fining-up trend. Cross-stratification with grading foresets. Well-defined internal erosional surfaces.	Bar deposits under subaerial sheet flows or stream flows.
<b>Sp.</b> Medium- to coarse-grained, cross-bedded sandstones (Fig. 7E)	Weakly sorted, medium- to coarse-grained sandstones which can contain scattered granules at the base.	Cross-stratification. Internal erosional surfaces. Few small scour structures.	Megaripple deposits under subaerial sheet flows or stream flows.
<b>Sr.</b> Massive, pedoturbated, well-sorted, fine-grained sandstones (Fig. 7D)	Massive, well-sorted, fine-grained sandstones.	Pedoturbation (root sleeves, destructuration of primary sedimentary	

**Table 1.** (continued).

Facies code. Facies name	Lithology	Sedimentary structures	Depositional processes
<b>Fm.</b> Massive, silty claystones with planar siltstone to fine-grained sandstone layers (Fig. 7D)	Reddish to yellowish (also purplish red in some horizons of plastic texture), massive, silty claystones with few thin (millimetric to centimetric) layers of well-sorted siltstones and very fine-grained sandstones.	structures by root growth). Locally few asymmetrical ripples at the top. Finning-up grading in siltstone to fine-grained sandstone layers.	Ripple deposits under subaerial sheet flows or stream flows. Deposits from suspension fallout in a floodplain. Overbank or waning flood deposits (siltstones and very fine-grained sandstones).
<b>P.</b> Pedogenetic carbonate nodules (Fig. 7F)	Indurated, cm to dm carbonate nodules, often scattered within plastic claystones, sometimes coalescent to form continuous carbonate horizons.	Vertical root casts.	Soil (Miall, 1978) or temporary vegetalized surface.

Gm, Gt, Sp, Sr, Fm and P are abbreviations from the facies code provided by Miall (1978).  
Gu and Sst are abbreviations defined by Barrier *et al.* (2010).

sequences, made up of alternate proximal and distal deposits, are arranged into overall trends measuring several tens of metres thick, recording progradational and retrogradational evolutions.

At the scale of the Guzargues basin, the large FA alternations are describing a whole progradational, then retrogradational, and finally progradational evolution, which is easily recognised on the field. These trends were used to correlate Priabonian deposits at basin scale and, combined with aerial photograph interpretation, to produce a detailed geological map that allows building cross-sections.


### 4.3 Tectonic-sedimentation relationships in the Guzargues basin

The detailed geological map (Fig. 7) and cross-sections (Fig. 8) of Guzargues basin show sequences of Priabonian conglomerates interfingering with marls, that are involved in gentle folding of complex geometry. To the south, the basin is separated from the contemporaneous Assas Basin by a NW-SE anticline, cored by Valanginian (Figs. 7 and 8 – section 1). North-south sections of Guzargues basin (Fig. 8 – sections 2 and 3) display an asymmetrical syncline structure: the southern limb is steeply dipping north, whereas the northern limb extends over 3 to 4 km in a gently dipping or subhorizontal position. In addition, the basal part of the Priabonian is affected by ENE-trending anticlines, while the youngest conglomeratic Priabonian seals the anticline (Fig. 8 – section 3).


*The Lirou anticline* – The Bartonian and the base of the Priabonian sequence (isolated sandstone and conglomerate beds in marls) are wedging towards an anticline with an EW-oriented, W-plunging axis. Conglomerate beds isolated in the silt and marls are observed on-lapping both flanks of the anticline (Fig. 6), and they display diverging bedding away

from the anticline (Figs. 7 and 9a). The upper conglomeratic interval, corresponding to the final progradational trend recorded by Priabonian deposits, progrades northwards across the whole basin, over a silt and shale interval and seals the Lirou anticline (Fig. 6). Folding affects the Lutetian lacustrine limestones, and the top of the Lutetian is eroded in the crest of the anticline. Variable thickness of Latest Cretaceous to Early Eocene marls suggests a decollement with the underlying monocline Valanginian limestones. The latter is affected by a sub-vertical, NE-trending fault, which displays a left-lateral component of movement (Fig. 8 – section 3). These observations lead to the following points: i) folding was initiated during Pyrenean shortening, and was still active during deposition of the early Priabonian; ii) folding of the Paleogene sequences is detached above the Valanginian limestone, through a detachment within the Late Cretaceous-Early Eocene marls; iii) folding is related with a NE-trending left lateral strike-slip.

*The southeastern boundary* – Figure 10 shows up to three stacked, angular unconformities, which affect the lower Priabonian sequences along the southern flank of an EW syn-sedimentary syncline. These angular unconformities pass northwards to correlative conformities, while they converge southwards, towards the basin boundary, corresponding to an E-trending anticline of Lutetian limestone. This anticline is locally eroded by a present-day transverse valley, but mapping reveals that sub-horizontal Priabonian conglomerates are exposed within this valley and that they laterally onlap onto the incised carbonates (Fig. 7). Clasts imbrication within the fluvial conglomerates indicates a consistent northwards paleocurrent in the paleo-valley. Very thin (several tens of metres) in the transverse valley, the Priabonian formation thickens rapidly northward within 1 kilometre distance, from less than 300 m to 450 m, and it exceeds 600 m in the syncline



**Fig. 5.** Photographs of outcrops illustrating some typical facies observed in the Priabonian deposits of the Guzargues basin. A: Facies Gu with carbonate clast alignment underlining tenuous cross-beddings. Note scours at the base of the channel (arrows). B: Facies Gt showing progressive transition between (base) unsorted, highly heterometric massive, stratified conglomerates, and (top) weakly sorted, cross-bedded conglomerates. C: Alternations between (Sst) cross-stratified sandstones with lenses of scattered subrounded gravels and pebbles, and (Gm) matrix-rich, graded, stratified conglomerates with weakly sorted pebbles mainly made of Mesozoic carbonates. D: Reddish to yellowish laminated silty claystones (facies Fm) with intercalation of a decimetric layer of massive fine-grained sandstones (facies Sr). E: Medium- to coarse-grained cross-bedded sandstones typical of facies Sp. F: 50 cm-thick carbonated horizon showing well-preserved, vertical root casts, probably derived from vegetalation on top of massive fine-grained sandstones (Sr). Note the occurrence of several pedogenetic carbonate nodules above the irregular top surface of the carbonated sandy horizon.




**Fig. 6.** Detailed stratigraphic correlation of Priabonian deposits along a 2 km-long N-S transect at the southeastern border of the Guzargues basin, showing differential subsidence and onlap geometries induced by the syndepositional growth of the Lirou anticline. See Figure 7 for location of logs. Horizontal correlation lines set at the maximum of progradation recorded by several meters-thick, stacked conglomeratic channels that closely rests on Lutetian limestones at the apex of the Lirou anticline.

axis. In addition, fluvial conglomerate belts pass laterally and distally (northwards) to silts and marls of alluvial plain facies. These observations clearly show that: i) the syncline was formed during deposition of the Priabonian; ii) the basin was fed by a north-flowing paleoriver, incised into the growing Lutetian limestone E-W anticline.

*Serre de Moujes* – The northern part of the basin corresponds to a syncline trending N020°, that involves Lutetian limestones and Priabonian conglomerates (Fig. 7). The conglomerates are more developed on the steep western flank, where they unconformably rest over the steeply dipping Lutetian limestones, and they grade eastwards to distal silts and marls (Fig. 8 – section 4). Along the eastern flank of the syncline, Priabonian formations are onlapping onto the Lutetian limestone; in addition, the Bartonian oncolith formation is transgressed towards the north by the Priabonian. Onlapping surfaces point to a pre-existing synform of Lutetian limestones prior to Priabonian deposition. The conglomerate belt is separated from that of the main part of the basin, suggesting the presence of a smaller, northern, sub-basin with a distinct feeder that supplies clastic sediments across the western boundary (Fig. 8 – section 2). Unfortunately, no clear evidence of paleocurrent was found. North-south correlations

across the entire Guzargues basins show that the northern sub-basin represents the youngest depocentre of the whole basin (Fig. 8 – section 1). To the NW, Priabonian is directly in contact with the Mesozoic carbonates through a sub-vertical faulted boundary. In the *Serre de Moujes*, outcrop in the damage zone of the bordering fault reveals sets of associated fractures and dissolution surfaces, consistent with a left-lateral strike-slip motion (Fig. 9b). Unfortunately, no slickenside could be observed to constrain the respective vertical and horizontal components of slip. *Saint-Bauzille-de-Montmel* borehole (located 1.5 km north, along strike) constrains a curved geometry for this fault and provides evidence for a 700 m vertical offset (Fig. 8 – section 4). Considering the fault geometry and the kinematic observations on the outcrop, a dominantly strike-slip movement along this bordering fault is suggested during Priabonian and the important vertical offset implies several kilometres of cumulated lateral offset. A significant part of such offset may be inherited from earlier (Pyrenean) tectonic phases. These observations argue for i) deposition of Priabonian syn-tectonic sediments in a hanging-wall syncline, controlled by the activity of the N020°-trending bounding fault, and ii) accommodation of a significant amount a left-lateral strike-slip, by this inherited fault.



**Fig. 7.** Detailed geological and structural map of the Guzargues basin (north of Highway D68, while Assas sub-basin is located south of the highway). Location of the sedimentological logs 1 to 9 reported in Figure 6 is indicated by fine black rectangles. Position of the structural cross-sections extremities (Figs. 8 and 9) is given by black arrows.

#### 4.4 Tectonics–paleoenvironment relationships in the Guzargues basin

Figure 11a provides a synthetic view of the paleoenvironment distribution across the Guzargues Basin during Priabonian. Fluvial sediments were delivered through an incised

canyon across a growing anticline, which separates Assas (to the south) and Guzargues basins. The belt of northward-flowing fluvial channels preferably occupied the most subsiding parts of the basin: i) in the south, parallel to the bordering growing anticline and, ii) in the overall N-trending axis of the syn-depositional syncline. The latter was controlled





Fig. 8. Structural sections across the Guzargues basin; see location in Figure 7.

by a set of NNE-trending faults which display map- and outcrop-scale evidences of normal faulting and left-lateral strike-slip. The secondary fluvial conglomerates depocenter of Serre de Moujes, in the northern part of the basin, witnesses an additional fluvial input to the basin, across a relay or a jog in the bordering fault system. The syn-depositional Priabonian

sequence is therefore controlled by folding of the Mesozoic through the Lutetian cover detached over the Triassic decollement, corresponding to an overall transcurrent, left-lateral kinematics, along the NNE-trending faults (Fig. 11b). The kinematics documented in the Guzargues basin result from NS shortening combined with EW extension.



**Fig. 9.** a: Internal onlap within the Priabonian conglomerates, which illustrates syndepositional folding of the Lirou anticline. Location on map [Figure 7](#). b: Penetrative deformation of the Jurassic limestones along the basin-bounding fault, indicating left-lateral kinematics. Serre de Moujes, location on map [Figure 7](#).

## 5 Les-Matelles and Saint-Martin-de-Londres Priabonian basins

### 5.1 Les-Matelles basin

The overall geometry of Les-Matelles basin (LMB) is that of an Oligocene syn-rift hanging-wall syncline, controlled by a NNE-trending normal fault ([Fig. 2](#)), known as the Matelles-Corconne Fault ([Benedicto \*et al.\*, 1999](#)). The Priabonian sequence mostly consists of marls and siltstones with rare conglomerate beds. Paleocurrent measurements in the channelized fluvial conglomerates indicate northwest to northeast flowing channels ([Egerton, 1996](#)). Detailed mapping ([Benedicto, 1996](#)) revealed that the underlying Priabonian sequence also presents growth structures in an asymmetric syncline, that displays a NW limb steeper ( $\geq 40^\circ$ ) and thinner than the SE limb ([Fig. 12](#)). Sequential tectonic restoration of transverse sections indicates an extensional offset across

Les-Matelles-Corconne Fault of some 375 m, during Priabonian ([Benedicto \*et al.\*, 1999](#)).


In the southernmost part of the basin, Priabonian unconformably overlies the syn-orogenic breccia of Paleocene age, related to the activity of the Montpellier thrust ([Andrieux \*et al.\*, 1971](#)). In the central segment of the basin, Priabonian is conformable over the distal syn-orogenic marls and oncoliths of Bartonian age. Finally, in the northern extension of this basin, Priabonian is unconformable over Neocomian marly limestones, which are deformed by EW-trending compressional structures. Such folds and thrusts are correlated to the Pic-Saint-Loup thrust, active during Bartonian ([Philip \*et al.\*, 1978](#)). Considered at the scale of the basin, the Priabonian is thus unconformable over pre-Pyrenean deformed series, as well as syn-orogenic Pyrenean sedimentation. Finally, the syn-rift Late Rupelian breccia is unconformably overlying the Priabonian ([Crochet, 1984](#)).

### 5.2 Saint-Martin-de-Londres basin


Saint-Martin-de-Londres basin (SMLB) extends in an EW asymmetric syncline, immediately north of the Pic-Saint-Loup thrust, a structure of the Pyrenean fold and thrust belt ([Fig. 2](#)). Above an unconformity over the Neocomian, the basin sequence starts with continental marls and thin beds of lacustrine limestones of latest Cretaceous to Paleocene age ([Crochet, 1984](#); [Freytet, 1971](#); [Philip \*et al.\*, 1978](#)), which are the distal correlative of the syntectonic “Vitrollian” breccia, deposited along the Montpellier thrust.

Lutetian lacustrine limestones, transgressive over Late Jurassic to Paleocene formations, form an 80 m to 100 m-thick slab that designs a wide, low amplitude, EW-oriented syncline ([Fig. 13](#)). The overlying breccia, dated to the Bartonian ([Philip \*et al.\*, 1978](#)), consists of local, late Jurassic, Neocomian and Lutetian limestones, derived from erosion of the active Pic-Saint-Loup thrust front. They pass northwards to distal continental marls and silts, which present syn-tectonic growth structures. The southern boundary of the basin has been intensely folded and faulted in relation with the Pyrenean Pic-Saint-Loup thrust, during Bartonian time ([Fig. 14](#)). The Priabonian sequence unconformably overlies the Bartonian breccia; to the east, it seals an intra-Bartonian thrust propagation fold ([Fig. 13](#)), which requires erosion of the Pyrenean-related structure prior to Priabonian deposition.

Along the southern margin of the SMLB, the Priabonian sequence designs an asymmetric syncline with sub-vertical to overturned southern limb ([Figs. 15a and 15b](#)). The sequence starts with a thin level of finely laminated limestone that contains bioclasts and charophytes characterising a lacustrine/palustrine environment. Laterally, the limestone passes to thicker and more massive beds of lacustrine mudstone. The thin basal limestone is correlated to the north of the basin, with 10 m to 15 m thick marls and carbonates deposited in an arid, evaporitic environment. Above the limestones, along the central southern margin only, are found grey marls including several thin beds of glauconitic sandstones with bioclasts and rare foraminifera ([Egerton, 1996](#)). This shallow marine sandstone passes upwards to marls and siltstones including thin bioturbated sandstone beds, deposited in a floodplain, similar to those described in LMB. In the upright, southern



**Fig. 10.** Detailed structural section in the south of Guzargues basin showing growth syncline and internal angular unconformities. See location of section 5 in Figure 7.



**Fig. 11.** a: Sketch map of the paleoenvironments in the Guzargues basin, in relation with the Priabonian tectonics. b: Conceptual tridimensional representation (no scale intended) of the interaction between Priabonian kinematics and sedimentation in the Guzargues basin. Braided fluvial conglomerates accumulate in the most subsiding parts, *i.e.* the actively deforming synclines that affect the Mesozoic-Lutetian cover, detached over the basement. See text for explanations.


limb of the syncline, these sandstone beds are affected by small scale reverse faults (Fig. 15c). They also display soft penetrative deformation of initially cylindrical root traces (Fig. 15d), which indicate an early, pre-lithification, N-S shortening prior to folding. Finally, up to 50 m stacked, channelized conglomerates, are deposited in several-metres-thick levels, interbedded with siltstones and marls. The high ratio of channelized conglomerates vs floodplain marls and siltstones is more in line with Guzargues than with Les-Matelles basins. Conglomerates are clast-supported and rather well-sorted. Clasts are polygenic, including significant number of quartz veins and black cherts, derived from the Paleozoic deformed basement. Distinctive clasts of glauconitic sandstones and calcarenites containing Albian orbitolines cannot be related to any local source, as the Mesozoic sequence is interrupted above the Valanginian due to the regional erosion of the Durancian Isthmus (Husson, 2013). Such clasts, also

been identified in Les-Matelles and Montferrier Priabonian outcrops, are imbricated under the influence of north to northwestwards paleocurrents (Egerton, 1996), and thus correspond to a southerly derived material (Freytet, 1971).


The Priabonian sequence of SMLB unconformable over Pyrenean structures, and containing clasts supplied by a southerly derived sedimentation was deposited in a N-S compressional setting, related to reactivation of the Pyrenean Pic-Saint-Loup Thrust.

## 6 Seismic sequence analysis of the Hérault Basin


The Hérault Basin (HB) is a NNE-SSW asymmetric half-graben developed along the Cévennes Fault, formed in relation with the rifting of the Gulf of Lion (Arthaud *et al.*, 1981;



**Fig. 12.** Structural section across Les-Matelles basin (from surface data and seismic profile) showing syn-extension deposition of Priabonian units. Modified after (Benedicto *et al.*, 1999).



**Fig. 13.** Detailed geological map of Saint-Martin-de-Londres basin. Position of sections in Figure 14 indicated as white line. The black dot and cross points to the churches of the villages.



**Fig. 14.** Structural section across St Martin-de-Londres basin: a: general; b: detailed, showing syn-depositional compression during Priabonian.

Maerten and Séranne, 1995; Séranne, 1999). It covers the western part of the Pyrenean Montpellier thrust (Fig. 2) and is filled by up to 1500 m-thick succession of rift and post-rift sequences. The syn-rift and earlier Pyrenean structures are therefore mostly covered, which lead us to investigate the Pyrenean to syn-rift transition through subsurface data.

The basin sequence consists of (i) Rupelian to Aquitanian, continental and lacustrine formations, and (ii) Burdigalian to Langhian, marine to fluvial post-rift deposits, forming syn-rift and post-rift sequences separated by a major unconformity. The pre-rift substratum consists of Permian to Jurassic sequence, as well as mid-Cretaceous bauxite deposits, which have been sampled by several boreholes (Lajoinie and Laville, 1979; Berger *et al.*, 1981). In addition, syn-tectonic Pyrenean Paleocene to Eocene formations (poorly exposed along the eastern margin of the basin) and Mesozoic carbonates deformed during Pyrenean compression can be extrapolated beneath the basin, and several Pyrenean thrusts repeating parts of the Jurassic sequence are documented (Berger *et al.*, 1981; Alabouvette *et al.*, 1982).

Two geophysical surveys acquired in 1983 and 1984, then reprocessed in 2005–2008 (Serrano and Hanot, 2005), provide a regular seismic coverage (Fig. 2) of sufficient quality to examine the seismic units underlying the syn-rift sequence (Fig. 16). Some of these seismic sections were already interpreted by Maerten and Séranne (1995) who focused their study on the geometry of syn- and post-rift units.


Reinterpretation of the seismic profiles allows us to identify, in several parts of the basin, a distinctive seismic unit below the syn-rift deposits (Fig. 16). It is bounded by angular unconformities both at the top and at the base and displays significant thickness variations and complex internal geometries. Geometrical relationships with underlying and overlying seismic sequences define its relative stratigraphy: (i) It is truncated by typical syn-rift sequences and is affected by the extensional syn-rift faults; (ii) It unconformably rests on pre-rift units affected by Pyrenean reverse faults (Fig. 17). This intermediate seismic unit, undrilled by any boreholes, does not correlate with surface exposures. However, comparison with

field observations in the GB, SMLB and LMB point out a similar structural and stratigraphic position suggestive of a Priabonian age. Mapping of this intermediate interval, of proposed Priabonian age, across the Hérault basin shows a discontinuous sequence including two main depocentres (Fig. 18a). One depocentre is located where the Cévennes Fault displays a significant change from an overall NE-SE direction to a SSE-trending orientation. H84X profile (Fig. 16) shows evidence for Priabonian syn-depositional, dip-slip movement across the SSE-trending fault segment. Such a geometry suggests a left-lateral movement along the Cévennes fault, associated with a dip-slip across the SSE segment which represents a releasing bend (*e.g.* (Christie-Blick and Biddle, 1985; Cunningham and Mann, 2007) of the master fault. A second depocentre is located along the SE margin of the Hérault basin, controlled by NE-trending faults that offset the earlier Pyrenean Montpellier thrust.


Isobaths of the base syn-rift (Fig. 18b) depict the depocentre of the syn- and post-rift Oligo-Miocene basin. Comparing the two maps (Figs. 18a and 18b) clearly shows that the successive sequences were controlled by two distinct tectonics settings and kinematics. The distribution and NE-SW overall orientation of the Oligo-Miocene depocentre, and the presence of a transfer zone (roughly parallel to H83E profile) are consistent with NW-SE extension across the Cévennes-fault (Maerten and Séranne, 1995). Our data do not support such a transfer zone active during the Priabonian.

## 7 Seismic sequence analysis in the Gardon-Saint-Chaptes basin

The Priabonian Gardon-Saint-Chaptes basin (GSCB) sits at the junction of an almost continuous NE-trending strip of Priabonian outcrops from the foreland of the Montpellier thrust (Fig. 2), to the large syn-rift Alès basin, parallel to the NE-trending Cévennes Fault. Syn-rift Oligocene sediments are observed, unconformable over the Priabonian, in the Alès Basins (Sanchis and Séranne, 2000) and in a distinct outcrop in



**Fig. 15.** a and b: General view of the eastern end of St-Martin-de-Londres Priabonian basin, an asymmetrical growth syncline, unconformable over the synorogenic Bartonian breccia, which were generated along the Pyrenean Pic-St-Loup thrust. c: Overturned Priabonian sandstone along the southern limb of the asymmetric syncline. d: Detail of the previous outcrop displaying small scale reverse faults that provide evidence for N-S compressive deformation. e: Deformed root traces, evidence for pre-lithification N-S compressive deformation of the Priabonian sandstone bed, prior to its folding. Outcrop location in [Figure 13](#).



**Fig. 16.** Seismic reflection profiles representative of the Hérault basin (H84D, H84F, H84X). The purple sequence interpreted as Priabonian (see discussion in text) is intermediate between the Pyrenean synorogenic units and the Oligocene syn-rift sequence. See location in Figures 2 and 18.


the southern part of the GSCB (Lettéron *et al.*, 2018). Transgression of the post-rift Burdigalian tidal sequences, restricted to the E-W paleo-ria of Uzès basin (Reynaud *et al.*, 2006), did not reach the Alès basin. The Gardon segment, between Alès and Saint-Chaptes, consists of a large asymmetrical syncline, with a distinctive NW-trend, contrasting with the general NE-trend of the contemporaneous basins (Fig. 2).

Pyrenean structures in this area consist of a succession of E-W folds affecting the thick Triassic to Neocomian sequence (Arthaud and Laurent, 1995), dated with syn-tectonic Paleocene to Bartonien terrigenous sediments in the syncline axes. The later are buried to the west beneath Priabonian and Oligocene sequences, which provides a first-order evidence for a major post-Pyrenean discontinuity (Fig. 2). Detailed mapping of the northeastern border of the GSCB also defines a marked angular unconformity of the Priabonian over the syn-Pyrenean breccia sequences (Sanchis, 2000).

Chronostratigraphy is rather well constrained due to numerous mammalian fossils (Fig. 3) reviewed in Lettéron *et al.* (2018). Priabonian deposits form a wide asymmetric

syncline, including a thick NE and reduced SW limbs, respectively. Seismic reflection survey of the Alès basin includes a southern profile (85LGC11), which reveals the structure and mode of formation of this syncline (Sanchis, 2000) (Fig. 19). In section, the GSCB displays Priabonian growth syncline migrating WNW-ward, which characterises dip-slip movement of a hanging-wall flat made of Mesozoic sequence detached above an eastward, low-dipping, decollement, above the Paleozoic basement. Such low-dipping decollement, the “Alès Fault”, has been identified in the neighbouring Alès basin and beneath the Mesozoic sequence, folded during the Pyrenean (Benedicto, 1996; Sanchis, 2000; Sanchis and Séranne, 2000). This decollement at the basement-cover interface is distinct from the Cévennes basement fault, although both the Cévennes and Alès faults emerge along the same trace. This seismic profile (Fig. 19) also provides evidence for 4 km E-SE migration of the depocentre, therefore suggesting similar displacement of the hanging-wall across the Alès Fault, in this direction, during Priabonian.

The Saint-Chaptes part of the GSCB has been investigated by seismic reflection profiling, aiming at hydrogeological exploration of the covered Barremian *Urgonian* karst (Josnin,



**Fig. 17.** Line drawings of parts of selected seismic profiles in the Hérault basin showing the lower and upper erosional boundaries of the intermediate sequence. See location in Figure 18.

2001). The sections tied to 4 wells sampling the Oligocene and Priabonian interval (Fig. 2) (Baral, 2018; Baral *et al.*, 2018) and correlated with surface exposures providing sedimentology and stratigraphy controls (Lettéron *et al.*, 2018), allowed to identify, date and map the major seismic markers across the basin (Fig. 20). In borehole, the Priabonian sequence consists of lacustrine marls and carbonates, that pass upward to sandstones. It rests on a thin interval of undated detrital continental sediment, which can be tentatively correlated with the syn-tectonic Bartonian observed in outcrops north of the borehole. Priabonian displays growth structures generated by extensional faults which ramp down into the Barremian (*Urgonian* facies) massive limestones. Such faults belong to the N-NE trending fault-network that truncates the Pyrenean folds, and is sealed by the Rupelian syn-rift sequence (Fig. 2). The dip-slip component of such faults observed in seismic, is not associated with any significant horizontal displacement, since the Pyrenean fold axes (Berger *et al.*, 1978) are not laterally offset (Fig. 2). Time-depth conversion was achieved using velocities stack provided with the seismic profiles and controlled with the borehole data (Baral, 2018), in order to make an isopach maps of the Priabonian depocentre (Fig. 21a). It provides evidence for Priabonian tectonics across N-NE-trending active faults, observed north of the basin, and southward, parallel to the Sommières basin (Fig. 2). The Gardon segment of the depocentre, although less well constrained by subsurface data, designs the NNW-trending asymmetrical syncline imaged in seismic (Fig. 19). The overlying Oligocene (syn-rift) depocentre (Fig. 21b) is restricted to the Saint-Chartes part of the GSCB and it is not affected by the earlier faults, nor controlled by any active boundary faults. It designs a wide depocentre, whose


subsidence axis strikes N050, oblique with respect with the Priabonian depocentre. This suggests a Rupelian syn-rift basin formed as a hanging-wall syncline (Benedicto *et al.*, 1999) above a detachment, which we interpret to be the Paleozoic basement-Mesozoic cover interface, *i.e.* the Alès Fault in this area. The structural relationships in the GSCB and the contrasted orientation of the Priabonian and Rupelian depocentres indicate two distinct events with different extension directions.

## 8 Regional synthesis and discussion

Field observations show that the GB, LMB and SMLB display similar Paleogene successions and stratigraphic settings. In all exposed basins, the syn-tectonic Pyrenean continental formations consist of breccia along the active thrusts that pass distally, *i.e.* northwards, to fine continental silts and marls, or lacustrine/palustrine limestones. They are overlain by Priabonian continental sediments, which were deposited in alluvial floodplains, including channelized polygenic, sub-rounded, conglomerates. Proximal-distal facies distributions, paleocurrents and some distinctive clast lithologies are consistent with a southern source for the clastic sediments. This is related with a remaining Pyrenean topography that extended in the present-day Gulf of Lion, until the Priabonian. Priabonian formations are preserved in distinct basins or sub-basins corresponding to subsiding depocentres, which display significant thickness variations and syn-tectonic growth structures. The lower boundary is an unconformity over the Pyrenean syn-orogenic formations and structures, although it can locally be observed conformable at outcrop scale. The angular unconformity is better expressed where Pyrenean tectonics was more developed: *i.e.* close to the Montpellier and Pic-Saint-Loup thrusts and around the anticlines. Priabonian sequences are eroded and unconformably overlain by Late Rupelian continental alluvial fans and lacustrine deposits, in relation with the rifting of the Gulf of Lion margin, except in the SMLB, where the stratigraphic record ends with Priabonian. The Priabonian sequences of the GSCB, observed in seismic profile and tied to borehole data, and those from the HB (although not precisely dated) provide evidence for similar structural and stratigraphic relationships, as observed in the field. Pyrenean structures and syn-orogenic sequences are unconformably overlain by a seismic sequence of highly variable thickness, which is also truncated by younger sequences displaying syn-rift growth structures against the basin bounding normal faults. We correlate this intermediate seismic sequence with the Priabonian interval described in the field in GB, LMB and SMLB.

At variance with the observed similar structural relationship with underlying and overlying sequences, the Priabonian deposits display contrasting structures in the different basins investigated (Fig. 22):

- LMB: Syn-tectonic growth-structures are consistent with a hanging-wall syncline, resulting from extensional faulting along the NNE-striking bounding normal fault.
- SMLB: Structures are related to syn-depositional compressional folding across the E-trending Pic-Saint-Loup thrust.




**Fig. 18.** Isopach maps of Priabonian (a) and of syn-rift (b) sequences in the Hérault basin, superimposed onto a simplified geological map (same legend as in Fig. 2, and location in Fig. 2). Contouring of depocenters in metres. The seismic profiles used in the study are represented in stippled green lines; plain green lines with labels correspond to the profiles shown in Figures 16 and 17. Faults are represented in red.

- GB: The tectonic sketch derived from the detailed geological map Figure 7 displays syn-depositional folding, which characterises both NE-SW compression and EW extension; in addition, there is evidence of left-lateral strike-slip along NNE-trending faults (Fig. 11b).
- HB: Depocentre distribution with respect with controlling faults suggests left-lateral motion along the NNE-trending Cévennes Fault, which accommodates normal faulting across N-trending splays (Fig. 22).
- GSCB: The Priabonian sequence was deposited as a hanging-wall syncline controlled by a SE-trending extensional low-angle decollement of the Mesozoic cover over the Paleozoic basement.


Such apparently contradicting kinematics can be understood in an overall left-lateral deformation along the regional NNE-trending faults, such as the Cévennes, Les-Matelles-Corconne and the Nîmes faults (Fig. 22). When submitted to a stress regime characterised by a horizontal, N-S, maximum principal stress  $\sigma_1$  and a horizontal E-W minimum principal stress  $\sigma_3$ , inherited faults oriented NNE are reactivated as left-lateral strike-slip faults, with releasing bends (Christie-Blick and Biddle, 1985) along more northerly oriented segments, as in LMB, HB, GSCB. East-west oriented fault splays such as the Pic-Saint-Loup correspond to restraining bends.

Syn-tectonic deposition in the syncline north of Uzès (Figs. 2 and 22) also provides additional evidence for N-S shortening across EW oriented structures, during Priabonian.

The amount of left-lateral displacement during Priabonian is poorly constrained, as it reactivates, and is superimposed, onto Pyrenean left-lateral ramps, resulting in a combined finite strain, which hampers the distinction of the two contributions to the strike-slip offset. In addition, our study shows that i) the strain was distributed across the whole area between the Cévennes and Nîmes faults and ii) possible discrepancies between the amount of deformation recorded in the Mesozoic cover and the basement may occur. However, the Priabonian tectonic-sedimentation relationships recorded in some of the studied basins provide minimum values for the offset. The extensional growth structure in the Hérault basin (Fig. 16c) suggests a minimum extension of some 3 km, corresponding to the minimum value of Priabonian left-lateral offset across this segment of the Cévennes fault. The migration of the Priabonian hanging-wall syncline in the GSCB (Fig. 19) indicates NE-ward displacement of 4 km. Restoration of the LMB points to 375 m Priabonian extension (Benedicto *et al.*, 1999) across a releasing bend of the Matelles-Corconne Fault. This could represent the order of magnitude of displacement across the other identified NE-trending faults that bound the Guzargues and Sommières basins (Fig. 2). We thus suggest



**Fig. 19.** Interpreted line drawing of a seismic reflection profile across the Gardon Basin showing the syn-depositional migration of the hanging-wall syncline. Modified after Sanchis (2000). Main markers are identified; Priabonian interval in purple overlay; intra-Priabonian marker corresponds to the base of “Grès de Celas” Formation. Location in Figures 2 and 21.




**Fig. 20.** Seismic reflection profile and corresponding line drawing in the Gardon-Saint-Chaptes basin showing evidence for syn-depositional (Priabonian) extensional faulting sealed by the Rupelian sequence. Maisonnnette borehole allows to interpret and to date the sequence. Modified after (Baral, 2018). Location in Figures 2 and 21.

that the Priabonian cumulated left-lateral motion distributed across the 40 km wide NE-trending wrenching zone remained moderate, with a minimal value of 4 to 5 kilometres. Even though some wrenching may have occurred without any syntectonic deposits recorded, the cumulated Priabonian movement unlikely exceeded 10 km. There is no data to document the offset across the major Nîmes fault.

The transition from Pyrenean compressional mountain building to rifting of the Gulf of Lion in Languedoc was

achieved by a specific evolution of the stress regime (Fig. 23). From latest Cretaceous to Bartonian, the N-S convergence of Sardinia and associated continental block with southern Europe were accommodated by E-W folds and thrusts, while the inherited regional NE-trending faults were activated as lateral-ramps. Such deformation results from a horizontal  $\sigma_1$  oriented N-S and a vertical  $\sigma_3$  (Arthaud and Laurent, 1995). During Priabonian, the stress regime evolved by permutation of  $\sigma_2$  and  $\sigma_3$  axes,  $\sigma_1$  remaining horizontal and oriented N-S.




**Fig. 21.** Isopach maps of Priabonian (a) and Rupelian (b) sequences in the Gardon-Saint-Chaptes basin. Note the different orientation of the Rupelian depocentre axis with respect to the previous Priabonian subsidence axis. Location in [Figure 2](#).


Such stress regime activated the inherited NE-trending faults as left-lateral strike-slip faults. This fault network, inherited from the Late Cretaceous-Bartonian Pyrenean orogeny, included restraining and releasing relays. The later corresponds to Priabonian depocentres, where  $\sigma_1$  and  $\sigma_2$  permuted. The restraining bend, where compression occurred ( $\sigma_1$  N-S horizontal and  $\sigma_3$  vertical), is illustrated by the Saint-Martin de Londres basin. Contemporaneous releasing bend and relay, where local extension occurred ( $\sigma_1$  vertical) is exemplified by the northern part of the Hérault basin, Les-Matelles basin, or Gardon-Saint-Chaptes basin ([Fig. 22](#)). Permanence of N-S horizontal  $\sigma_1$  from Late Cretaceous through to Late Eocene was also documented in southern Provence, defined by both strike-slip and reverse faults ([Lacombe \*et al.\*, 1992](#)). N-S shortening along the regional NE-trending faults accommodated transpression of the oblique Africa-Eurasia convergence in the Languedoc-Provence-Corsica domain ([Lacombe and Jolivet, 2005](#)) and lasted until Priabonian. During Late Rupelian, the stress regime rotated clockwise around the vertical  $\sigma_1$ , by  $30^\circ$  ([Séranne, 1999](#)), which activated the inherited NE-trending strike-slip faults as extensional faults. This Rupelian change in stress orientation reflects a change at plate boundaries, which we relate to the onset of subduction roll-back and back-arc rifting above the Apennine subduction ([Réhault \*et al.\*, 1984](#); [Séranne, 1999](#); [Jolivet and Faccenna, 2000](#); [Jolivet \*et al.\*, 2015](#); [Romagny \*et al.\*, 2020](#)). Southeast-trending transfer faults zones are a distinctive feature of the rifting of the Gulf of Lion ([Gorini \*et al.\*, 1993](#); [Guennoc \*et al.\*, 2000](#); [Mauffret \*et al.\*, 2001](#); [Canva \*et al.\*, 2020](#); [Maillard \*et al.\*, 2020](#)). However, there is no evidence of active SE-oriented

structure in the onshore study area, until the Oligocene rifting episode ([Maerten and Séranne, 1995](#); [Sanchis and Séranne, 2000](#)). The Priabonian strike-slip deformation documented in the present study therefore constitutes a distinct tectonic phase, corresponding to different kinematics and driving forces than the rifting of the Gulf of Lion ([Séranne, 1999](#)). We suggest that the development of SE trending transfer zones in the back-arc basin, as from the Rupelian, dates the initiation of slab tearing in the underlying subduction zone ([Romagny \*et al.\*, 2020](#)).

Priabonian left-lateral wrenching in Languedoc links the active Pyrenean orogen to the E-W rifting of the European Cenozoic Rift System (ECRIS) ([Dèzes \*et al.\*, 2004](#)) and the North Alpine Foreland Basin ([Ford and Lickorish, 2004](#)). Such left-lateral wrenching during Priabonian most probably extended southward in the Corbières ([Fig. 1](#)); however, lack of Priabonian sedimentary record in this area does not allow to confirm this. Eocene northward motion of the Adriatic plate was accommodated by left-lateral oblique convergence along the SW Alps until continental collision in Oligocene ([Ford \*et al.\*, 2006](#); [Dumont \*et al.\*, 2012](#)) ([Fig. 24](#)). Convergence between the Iberia and Europe, respectively, lasted until early Miocene in the Pyrénées *e.g.* ([Mouthereau \*et al.\*, 2014](#); [Teixell \*et al.\*, 2018](#)) and *circa* 85 km shortening is computed for the Late Eocene to Early Oligocene interval ([Teixell \*et al.\*, 2016](#)). Pyrenees shortening in Corbières, Languedoc and Provence terminated during the Bartonian ([Séguret and Benedicto, 1999](#); [Alabouvette \*et al.\*, 2001](#); [Bestani \*et al.\*, 2015](#)). Northward flowing clastic sedimentation pattern in Languedoc, Provence and the SE Alps shows that this segment of the orogen, located




**Fig. 22.** Synthetic regional map of Priabonian structures and kinematics in Languedoc. For location of specific features and localities, please refer to [Figure 2](#), which shares the same frame. The study area corresponds to a zone of sinistral shearing between the Cévennes and Nîmes faults, with a minimal value of 5 km (grey overlay symbolises the simple shear).



**Fig. 23.** Evolution of stress regime from mid-Eocene Pyrenean shortening, through Priabonian strike-slip regime, to Rupelian rifting.

in the present-day Gulf of Lion, was being dismantled at that time (Guieu and Roussel, 1990; Egerton, 1996; Joseph and Lomas, 2004).

The left-lateral movement along the Nîmes and Cévennes fault system extended northward to the Jura region where sinistral strike-slip is well documented during Eocene (Homberg *et al.*, 2002). It is also consistent with the transfer zone between Rhine and Bresse grabens (Bergerat *et al.*, 1990; Lacombe *et al.*, 1993; Madritsch *et al.*, 2009), which both record a distinctive Priabonian pulse of sedimentation predating a mid-Oligocene extensional tectonics (Sissingh, 1998). Crustal extension across these rift basins amounts to 7 km (Ziegler and Dèzes, 2005), a value in agreement with the offset measured across the strike-slip faults.



**Fig. 24.** Simplified map Pyrénées/European Cenozoic Rift System/Alps during Priabonian. Paleogeographic reconstruction at 35 Ma from Romagny *et al.* (2020). Position of the North Alpine foreland from Ford and Lickorish (2004) and Dumont *et al.* (2012) and of the ECRIS from Dèzes *et al.* (2004).

During the Priabonian, the Languedoc-Provence segment of the Pyrenean orogen was thus surrounded by two active convergence zones, while it was undergoing collapse. This different behaviour may be due to the interaction with the west-European rifting (Dèzes *et al.*, 2004), which propagated southward beneath the Languedoc-Provence-Pyrénées.

## 9 Conclusion

The transition from the Pyrenean orogeny to the Gulf of Lion rifting in onshore Languedoc occurred during Priabonian. Prior to Priabonian time, every structural and tectonic-sedimentation evidence argue for Pyrenean compressional deformation. In contrast, after Priabonian, all structures indicate NW-SE extension related to the Gulf of Lion rifting. This makes the Priabonian interval a key intermediate period of kinematic change. Reassessment of the chronostratigraphic evolution of the area allowed us to constrain the calendar of events. This period is recorded by deposition of a distinctive sedimentary sequence, which is documented both by surface analyses and seismic profiling of Priabonian basins. Such intermediate sequence is bounded at the base by an angular unconformity over the Pyrenean structures, and at the top by the Oligocene syn-rift unconformity, respectively. Continental sedimentation ranges from lacustrine/palustrine limestone and evaporitic marls at the base, to marls and silts of alluvial plain, including variable proportion of sandstone sheets and channelized fluvial conglomerates, toward the top. The Priabonian series progrades north- or northeastward and is derived from a southern source, presently located offshore, which indicates the existence of a remaining Pyrenean topography in the present-day Gulf of Lion.

Field mapping and seismic analyses of the tectonic-sedimentation relationships in the Priabonian basins give evidence for syn-depositional deformation against bounding faults and in growth synclines. Contrasted modes of deformation are observed, within short distance, in the Priabonian basins: i) controlled by extensional movement of NNE to NNW-trending faults or segments of faults, or ii) controlled by E-W-oriented thrusts and folds. Regional mapping of these structures shows a distribution of the compressional and extensional structures which are consistent with a left-lateral shearing of the area comprised between the major, NE-trending, Cévennes and Nîmes faults. Such sinistral strike-slip of moderate offset ( $\pm 5$  km) accommodates the E-W extension of the West European rifts basins that are located to the north of the study area. To the south, the strike-slip faults are connected to the ongoing shortening in the Spanish Pyrénées. Priabonian deformation and sedimentation records the initial stages of the dismantling of the Languedoc-Provence segment of Pyrénées, located between two active orogens: the Spanish Pyrénées and the Western Alps. The late orogenic dismantling of the eastern Pyrénées (Jolivet *et al.*, 2020) therefore started up to 5 My earlier than the Rupelian onset of Apennine subduction retreat and associated back-arc NW-SE extension.

*Acknowledgements.* This study was carried out during the OROGEN project, funded by TOTAL, BRGM and CNRS-INSU. It benefited from the stimulating discussions amongst the OROGEN community.

We acknowledge the BRGM for providing the reprocessed lines of the H83 and H84 seismic surveys, presented in this study. We are indebted to SMAGE (*Syndicat Mixte pour l'Aménagement et la Gestion Equilibrée des Gardons*) for providing us with subsurface data for the Gardon-Saint-Chaptes basin, and for support for C.B.

We are extremely grateful to J.Y. Crochet, J.A. Remy, F. Lihoreau, and R. Tabuce (ISEM-University of Montpellier) for discussing palaeontology and mammal biostratigraphy, and for guiding us to the precise locations of ancient and more recent paleontological discoveries.

The paper benefited from fruitful exchanges with Flavia Girard and Renaud Caby (Géosciences Montpellier), François Fournier (CEREGE), Alexandre Letteron (IFP-CEREGE), Hubert Camus (Cenote) and Pascal Fénart (Hydrofisis).

The manuscript has been improved (and extended!) thanks to the insightful comments of François Guillocheau, Sylvain Calassou, and of an anonymous reviewer. The precious suggestions and corrections of the Guest Editor, Olivier Lacombe, are greatly appreciated.

## References

- Aguilar JP, Legendre S, Michaux J. 1997. In: *Actes du Congrès BiochroM'97-Montpellier, 14-17 avril 1997*, Montpellier, 818 p.
- Alabouvette B, Arthaud F, Feist R, Medioni R, Brousse R. 1982. Carte géologique de la France au 1/50 000. Feuille de Lodève. BRGM.
- Alabouvette B, Berger G, Demange M, Guérangé-Lozes J. 2001. Montpellier - Carte géologique de la France à 1/250 000<sup>e</sup>. BRGM.
- Andrieux J, Mattauer M, Tomasi P, Martinez C, Reille JL, Matte P, *et al.* 1971. Carte géologique de la France au 1/50 000. Feuille de Montpellier N°990. BRGM.

- Arthaud F, Laurent P. 1995. Contraintes, déformation et déplacement dans l'avant-pays Nord-pyrénéen du Languedoc méditerranéen. *Geodynamica Acta* 8: 142–157.
- Arthaud F, Séguret M. 1981. Les structures pyrénéennes du Languedoc et du Golfe du Lion (Sud de la France). *Bulletin de la Société géologique de France* XXIII(1): 51–63.
- Arthaud F, Mégard F, Séguret M. 1977. Cadre tectonique de quelques bassins sédimentaires. *Bull. Centres Explor. – Prod. Elf Aquitaine* 1 (1): 147–188.
- Arthaud F, Ogier M, Séguret M. 1981. Géologie et géophysique du Golfe du Lion et de sa bordure nord. *Bulletin du BRGM* 1(3): 175–193.
- Baral C. 2018. Caractérisation géologique du réservoir karstique profond du bassin de Saint-Chaptes, Gard, France. Master 2, Montpellier, 50 p.
- Baral C, Séranne M, Fénart P, Camus H. 2018. Geological characterization and formation of a covered karstic reservoir in the Urgonian limestones of the Saint Chaptes Basin (Gard, South of France): towards predicting the distribution and state of karstification. In: *Proceedings Eurokarst 2018, 2–5 juillet 2018*, Besançon, France.
- Barrier L, Proust J-N., Nalpas T, Robin C, Guillocheau F. 2010. Control of alluvial sedimentation at foreland-basin active margins: a case study from the Northeastern Ebro basin (Southeastern Pyrénées, Spain). *Journal of Sedimentary Research* 80: 728–749.
- Benedicto A. 1996. Modèles tectono-sédimentaires de bassins en extension et style structural de la marge passive du Golfe du Lion (Partie Nord), sud-est de la France. Université de Montpellier II, 235 p.
- Benedicto A, Labaume P, Séguret M, Séranne M. 1996. Low-angle crustal ramp and basin geometry in the Gulf of Lion passive margin: the Oligocene-Aquitainian Vistrenque graben, SE France. *Tectonics* 15(6): 1192–1212.
- Benedicto A, Séguret M, Labaume P. 1999. Interaction between faulting, drainage and sedimentation in extensional hanging-wall syncline basins: example of the Oligocene Matelles Basin, Gulf of Lion margin (S.E. France). In: Durand B, Jolivet L, Horváth F, Séranne M, eds. *The Mediterranean basins: Tertiary extension within the Alpine Orogen*. Volume Special Publication 156. London: The Geological Society, pp. 81–108.
- Berger G, Lemprière P, Séguret M. 1978. Carte géologique de la France au 1/50 000. Feuille d'Anduze n°938. BRGM.
- Berger G, Feist R, Freyret P. 1981. Carte géologique de la France au 1/50 000. Feuille de Pézenas. BRGM.
- Bergerat F, Mugnier J-L, Guellec S, Truffert C, Cazes M, Damotte B, *et al.* 1990. Extensional tectonics and subsidence of the Bresse basin: an interpretation from ECORS data. *Mémoire de la Société Géologique de France* 156: 145–156.
- Bestani L. 2015. Géométrie et cinématique de l'avant-pays provençal : modélisation par coupes équilibrées dans une zone à tectonique polyphasée. PhD Doctorat, Université Aix-Marseille, 244 p.
- Bestani L, Espurt N, Lamarche J, Floquet M, Philip J, Bellier O, *et al.* 2015. Structural style and evolution of the Pyrenean-Provence thrust belt, SE France. *Bulletin de la Société géologique de France* 186 (ILP: Lithosphere dynamics of sedimentary basins: The Circum-Mediterranean basins and analogues): 223–241.
- Bestani L, Espurt N, Lamarche J, Bellier O, Hollender F. 2016. Reconstruction of the Provence Chain evolution, southeastern France. *Tectonics* 35(6): 1506–1525.
- Canva A, Thion I, Peyrefitte A, Couëffé R, Maillard A, Jolivet L, *et al.* 2020. The Catalan magnetic anomaly: its significance for the crustal structure of the Gulf of Lion passive margin and relationship to the Catalan transfer zone. *Marine and Petroleum Geology* 113: 104174.
- Chelalou R, Nalpas T, Bousquet R, Prevost M, Lahfid A, Pujol M, *et al.* 2016. New sedimentological, structural and paleo-thermicity data in the Boucheville Basin (eastern North Pyrenean Zone, France). *Comptes Rendus Géosciences* 348 (From rifting to mountain building: the Pyrenean belt): 312–321.
- Chevrot S, Sylvander M, Diaz J, Martin R, Mouthereau F, Manatschal G, *et al.* 2018. The non-cylindrical crustal architecture of the Pyrénées. *Scientific Reports* 8: 9591.
- Christie-Blick N, Biddle KT. 1985. Deformation and basin formation along strike-slip. In: Christie-Blick N, Biddle KT, eds. *Strike-slip deformation, basin formation, and sedimentation*. Volume Special Publication 37. SEPM, pp. 1–34.
- Christophoul F, Ford M, Grool A, Rougier G, Hemmer L. 2016. Tectono-stratigraphic evolution of the northeastern Pyrenean Foreland. *Proceedings International Geological Congress* 35: 3188.
- Cochelin B, Lemirre B, Denèle Y, De Saint-Blanquat M, Lahfid A, Duchêne S. 2018. Structural inheritance in the Central Pyrénées: the Variscan to Alpine tectonometamorphic evolution of the Axial Zone. *Journal of the Geological Society of London* 175(2): 336–351.
- Crochet JY. 1984. Géologie et paléontologie de la partie septentrionale du fossé oligocène des Matelles (Hérault, sud de la France). *Géologie de la France* 1-2: 91–104.
- Crochet J-Y, Godinot M, Hartenberger J-L, Rémy JA, Sigé B, Sudre J. 1988. Découverte dans le bassin de Saint-Martin-de-Londres (Hérault, Sud de la France) d'un gisement à vertébrés continentaux d'âge Eocène moyen. *Cour. Forsch. – Inst. Senckenberg* 107: 419–434.
- Crochet J-Y, Hartenberger J-L, Rémy J-A, Sudre J, Welcome J-L. 1997. Découverte de vertébrés continentaux de l'Eocène moyen et supérieur dans le bassin des Matelles (Hérault, Sud de la France) et redécouverte du «*Lophiodon* des Matelles». *Géologie de la France* 1: 35–45.
- Cunningham WD, Mann P, eds. 2007. Tectonics of strike-slip restraining and releasing bends. Volume Special Publication 290. London: The Geological Society, pp. 1–12.
- Dautria J-M, Liotard J-M, Bosch D, Alard O. 2010. 160 Ma of sporadic basaltic activity on the Languedoc volcanic line (southern France): a peculiar case of lithosphere-asthenosphere interplay. *Lithos* 120: 202–222.
- Debrand-Passard S, Courbouleix S. 1984. Synthèse géologique du Sud-Est de la France – Stratigraphie et paléogéographie. Orléans, France: BRGM, 615 p.
- Debroas E-J. 1990. Le flysch noir albo-cénomaniem témoin de la structuration albienne à sénonienne de la Zone nord-pyréenne en Bigorre (Hautes-Pyrénées, France). *Bulletin de la Société géologique de France* VI(2): 273–285.
- Dèzes P, Schmid SM, Ziegler PA. 2004. Evolution of the European cenozoic rift system: interaction of the Alpine and Pyrenean orogens with their foreland lithosphere. *Tectonophysics* 389: 1–33.
- Dumont T, Schwartz S, Guillot S, Simon-Labrie T, Tricart P, Jourdan S. 2012. Structural and sedimentary records of the Oligocene revolution in the Western Alpine arc. *Journal of Geodynamics* 56-57: 18–38.
- Egerton PD. 1996. Inherited drainage pathways sediment distribution and provenance in rift basins. PhD, University of Leeds, 386 p.
- Feist-Castel M. 1971. Sur les Charophytes fossiles du bassin Tertiaire d'Alès (Gard). *Geobios* 4(3): 157–172.
- Ford M, Lickorish WH. 2004. Foreland basin evolution around the western Alpine arc. In: Joseph P, Lomas S, eds. *Deep-water sedimentation in the Alpine basin of SE France: New perspectives on the Grbs d'Annot and related systems*. Volume Special Publication 221. London: The Geological Society, pp. 39–63.
- Ford M, Vergès J. 2020. Evolution of a salt-rich transtensional rifted margin, eastern North Pyrénées, France. *Journal of the Geological Society* 178(1): 2019–2157.

- Ford M, Duchêne S, Gasquet D, Vanderhaeghe O. 2006. Two-phase orogenic convergence in the external and internal SW Alps. *Journal of the Geological Society of London* 163: 815–826.
- Freytet P. 1971. Les dépôts continentaux et marins du Crétacé supérieur et des couches de passage à l'Eocène en Languedoc. *Bulletin du BRGM* 4: 1–54.
- Gastaud J, Campredon R, Féraud G. 1983. Les systèmes filoniens des Causses et du Bas Languedoc (Sud de la France): géochronologie, relations avec les paléocontraintes. *Bulletin de la Société Géologique de France* XXV(5): 737–746.
- Gaviglio P, Gonzales J-F. 1987. Fracturation et histoire tectonique du bassin de Gardanne (Bouches-du-Rhône). *Bulletin de la Société Géologique de France* 8(4): 675–682.
- Gorini C, Viallard P, Deramond J. 1991. Modèle d'inversion négative: la tectonique extensive post-nappe du fossé de Narbonne-Sigean (Corbières, sud de la France). *C. R. Acad. Sci. Paris* 312: 1013–1019.
- Gorini C, Le Marrec A, Mauffret A. 1993. Contribution to the structural and sedimentary history of the Gulf of Lions (western Mediterranean), from the ECORS profiles, industrial seismic profiles and well data. *Bulletin de la Société géologique de France* 164(3): 353–363.
- Gorini C, Mauffret A, Guennoc P, Le Marrec A. 1994. Structure of the Gulf of Lions (Northwest Mediterranean Sea): a review. In: Mascle A, ed. *Hydrocarbon and petroleum geology of France*. Volume Special Publication of the European Association of Petroleum Geology 4. Berlin: Springer-Verlag, pp. 223–243.
- Grambast L. 1962. Aperçu des charophytes tertiaires du Languedoc et leur signification stratigraphique. *C. R. Somm. Soc. Géol. France* 10: 313–314.
- Guennoc P, Gorini C, Mauffret A. 2000. Histoire géologique du Golfe du Lion et cartographie du rift oligo-aquitainien et de la surface messinienne. *Géologie de la France* 3: 67–97.
- Guiu G, Roussel J. 1990. Arguments for the pre-rift uplift and rift propagation in the Ligurian-Provençal basin (Northwestern Mediterranean) in the light of Pyrenean Provençal orogeny. *Tectonics* 9(5): 1113–1142.
- Hampton MA, Horton BK. 2007. Sheetflow fluvial processes in rapidly subsiding basin, Altiplano plateau, Bolivia. *Sedimentology* 54: 1121–1147.
- Hartenberger J-L, Sigé B, Sudre J. 1969. Les gisements de vertébrés de la région montpellieraine–1. Gisements éocènes. *Bulletin BRGM* 1(1): 7–18.
- Hippolyte JC, Angelier J, Bergerat F, Nury D, Guiu G. 1993. Tectonic-stratigraphic record of paleostress time changes in the Oligocene basins of the Provence, southern France. *Tectonophysics* 226: 15–35.
- Homberg C, Bergerat F, Philippe Y, Lacombe O, Angelier J. 2002. Structural inheritance and cenozoic stress fields in the Jura fold-and-thrust belt (France). *Tectonophysics* 357: 137–158.
- Horton BK, Schmitt JG. 1996. Sedimentology of a lacustrine fan-delta system, Miocene Horse Camp Formation, Nevada, USA. *Sedimentology* 43: 133–155.
- Husson E. 2013. Interaction géodynamique/karstification et modélisation géologique 3D des massifs carbonatés: implication sur la distribution prévisionnelle de la karstification. Exemple des paléokarsts crétacés à néogènes du Languedoc montpelliérain. Doctorat, Université Montpellier 2, 314 p.
- Jolivet L, Faccenna C. 2000. Mediterranean extension and the Africa-Eurasia collision. *Tectonics* 19(6): 1095–1106.
- Jolivet L, Gorini C, Smit J, Leroy S. 2015. Continental breakup and the dynamics of rifting in back-arc basins: The Gulf of Lion margin. *Tectonics* 34. <https://doi.org/10.1002/2014TC003570>.
- Jolivet L, Romagny A, Gorini C, Maillard A, Thion I, Couëffé R, *et al.* 2020. Fast dismantling of a mountain belt by mantle flow: late-orogenic evolution of Pyrénées and Liguro-Provençal rifting. *Tectonophysics* 776: 228312.
- Joseph P, Lomas S, eds. 2004. Deep-water sedimentation in the Alpine Foreland Basin of SE France: New perspectives on the Grès d'Annot and related systems – An introduction. In: *Deep-water sedimentation in the Alpine basin of SE France: New perspectives on the Grès d'Annot and related systems*. Volume Special Publication 221. London: The Geological Society, pp. 1–16.
- Josnin JY. 2001. Influence de la tectonique quaternaire sur la structure de deux aquifères karstiques du Languedoc (France). *Eclogae geologicae Helveticae* 94: 173–184.
- Lacombe O, Jolivet L. 2005. Structural and kinematic relationships between Corsica and the Pyrénées-Provence domain at the time of the Pyrenean orogeny. *Tectonics* 24(1): TC1003.
- Lacombe O, Angelier J, Laurent P. 1992. Determining paleostress orientations from faults and calcite twins: a case study near the Sainte-Victoire Range (southern France). *Tectonophysics* 201 (1-2): 141–156.
- Lacombe O, Angelier J, Byrne D, Dupin JM. 1993. Eocene-Oligocene tectonics and kinematics of the Rhine-Saone continental transform zone (Eastern France). *Tectonics* 12(4): 874–888.
- Lagabrielle Y, Bodinier J-L. 2008. Submarine reworking of exhumed subcontinental mantle rocks: field evidence from the Lherz peridotites, French Pyrénées. *Terra Nova* 20(1): 11–21.
- Lagabrielle Y, Labaume P, de Saint Blanquat M. 2010. Mantle exhumation, crustal denudation, and gravity tectonics during Cretaceous rifting in the Pyrenean realm (SW Europe): insights from the geological setting of the lherzolite bodies. *Tectonics* 29: TC4012, 4011–4026.
- Lajoinie J-P, Laville P. 1979. Les formations bauxitiques de la Provence et du Languedoc. Dimensions et distribution des gisements. *Mémoire du BRGM* 100: 146.
- Lamotte DFD, Souque C, Grelaud S, Robion P. 2002. Early record of tectonic magnetic fabric during inversion of a sedimentary basin. Short review and examples from the Corbières transfer zone (France). *Bulletin de la Société géologique de France* 172(5): 461–469.
- Le Pichon X, Rangin C, Hamon Y, Loget N, Lin JY, Andreani L, *et al.* 2010. Geodynamics of the France Southeast Basin. *Bulletin de la Société géologique de France* 181(6): 477–501.
- Lettéron A, Hamon Y, Fournier F, Séranne M, Pellenard P, Joseph P. 2018. Reconstruction of a saline, lacustrine carbonate system (Priabonian, St-Chartes Basin, SE France): depositional models, paleogeographic and paleoclimatic implications. *Sedimentary Geology* 367: 20–47.
- Mack GH, Leeder MR. 1999. Climatic and tectonic controls on alluvial fan and axial fluvial sedimentation in the Plio-Pleistocene Palomas half graben, southern Rio Grande Rift. *Journal of Sedimentary Research* 69: 635–652.
- Madritsch H, Kounov A, Schmid SM, Fabbri O. 2009. Multiple fault reactivations within the intra-continental Rhine-Bresse Transfer Zone (La Serre Horst, eastern France). *Tectonophysics* 471: 297–318.
- Maerten L, Séranne M. 1995. Extensional tectonics of the Oligo-Miocene Hérault Basin (S France), Gulf of Lion margin. *Bulletin de la Société géologique de France* 6: 739–749.

- Maillard A, Jolivet L, Lofi J, Thion I, Couëffé R, Canva A, *et al.* 2020. Transfer zones and associated volcanic province in the eastern Valencia Basin: Evidence for a hot rifted margin? *Marine and Petroleum Geology* 119: 104419.
- Masclé A, Vially R. 1999. The petroleum systems of the South-East Basin and Gulf of Lions (France). In: Durand B, Jolivet L, Horváth F, Séranne M, eds. *The Mediterranean Basins: tertiary extension within the Alpine Orogen*. Volume Special Publication 156. London: The Geological Society, pp. 121–140.
- Mattauer M. 2002. La faille normale oligocène des Matelles (environs de Montpellier). Une difficile confrontation entre un modèle à la mode et les observations de terrain. *Géologie de la France* 1: 39–43.
- Mauffret A, Gorini C. 1996. Structural style and geodynamic evolution of Camargue and Western Provençal basin, southeastern France. *Tectonics* 15(2): 356–375.
- Mauffret A, Durand de Grossouvre B, Dos Reis AT, Gorini C, Nercissian A. 2001. Structural geometry in the eastern Pyrénées and western Gulf of Lion (Western Mediterranean). *Journal of Structural Geology* 23: 1701–1726.
- Miall AD. 1977. A review of the braided river depositional environment. *Earth-Science Reviews* 13: 1–62.
- Miall AD. 1978. Facies types and vertical profile models in braided river deposits: a summary. *Fluvial sedimentology*. Volume Memoir 5. Canadian Society of Petroleum Geologists, pp. 597–604.
- Mouthereau F, Filleaudeau P-Y, Vacherat A, Pik R, Lacombe O, Fellin MG, *et al.* 2014. Placing limits to shortening evolution in the Pyrénées: role of margin architecture and implications for the Iberia/Europe convergence. *Tectonics* 33: 2242–2314.
- Nemec W, Postma G. 1993. Quaternary alluvial fans in the southwestern Crete: Sedimentation processes and geomorphic evolution. In: Marzo M, Puigdefabregas C, eds. *Alluvial sedimentation*. Volume Special Publication 17. International Association of Sedimentologists, pp. 235–276.
- Philip H, Mattauer M, Bodeur Y, Séguret M, Puech JP, Mattei J. 1978. Carte géologique de la France au 1/50 000. Feuille de St Martin de Londres N°963. BRGM.
- Réhault JP, Boillot G, Mauffret A. 1984. The western Mediterranean basin geological evolution. *Marine Geology* 55: 447–477.
- Rémy J-A. 1985. Nouveaux gisements de mammifères et reptiles dans les Grès de Célas (Eocène sup. du Gard). *Palaeontographica* 189: 171–225.
- Remy J-A. 2015. Les Périssodactyles (Mammalia) du gisement Bartonien supérieur de Robiac (Éocène moyen du Gard, Sud de la France). *Palaeovertebrata* 23(1-4): 211–216.
- Remy J-A, Fournier F. 2003. Mammifères fossiles des grès de Célas (Eocène supérieur du Gard): découvertes récentes. *Bulletin de la Société d'Etudes des Sciences Naturelles de Nîmes et du Gard* 64: 18–30.
- Rémy J-A, Lesage J-L. 2005. Un nouveau gisement de vertébrés d'âge Priabonien et son contexte géologique (Tranchée de Nozières, Gard). *Bulletin de la Société d'Etudes des Sciences Naturelles de Nîmes et du Gard* 65: 7–15.
- Rey R. 1962. Sur l'importance paléontologique des faunes malacologiques continentales de très petites dimensions comme celles de la région de Montpellier. *C. R. Somm. Soc. Géol. France* 10: 312–313.
- Reynaud J-Y, Dalrymple RW, Vennin E, Parize O, Besson D, Rubino JL. 2006. Topographic controls on production and deposition of tidal cool-water carbonates, Uzès Basin, SE France. *Journal of Sedimentary Research* 76: 116–129.
- Romagny A, Jolivet L, Menant A, Bessière E, Maillard A, Canva A, *et al.* 2020. Detailed tectonic reconstructions of the Western Mediterranean region for the last 35 Ma: insights on driving mechanisms. *Bulletin de la Société géologique de France* 191: art n°37.
- Sanchis E. 2000. Méthodologie d'imagerie 3D des bassins : des exemples dans le bassin du Sud Est. BRGM/Univ-Montpellier 2, 70 p.
- Sanchis E, Séranne M. 2000. Structural style and tectonic evolution of a polyphase extensional basin of the Gulf of Lion passive margin: the Tertiary Alès Basin, southern France. *Tectonophysics* 322: 243–264.
- Séguret M, Benedicto A. 1999. Le duplex à plis de propagation de rampes de Cazedarnes (arc de Saint-Chinian, avant-pays nord-pyrénéen, France). *Bulletin de la Société géologique de France* 170 (1): 31–44.
- Séguret M, Benedicto A, Séranne M. Structure profonde de la région du Gard Rhodanien – Apport du retraitement et de la réinterprétation de données sismiques régionales. In: *Proceedings Journées Scientifiques Gard, Bagnols/Cèze (Gard)*, Volume Atlas des posters, 1997, pp. 31–32.
- Séranne M. 1999. The Gulf of Lion continental margin (NW Mediterranean) revisited by IBS: an overview. In: Durand B, Jolivet L, Horváth F, Séranne M, eds. *The Mediterranean basins: Tertiary extension within the Alpine Orogen*. Volume Special Publication 156. London: The Geological Society, pp. 15–36.
- Séranne M, Benedicto A, Truffert C, Pascal G, Labaume P. 1995. Structural style and evolution of the Gulf of Lion Oligo-Miocene rifting: Role of the Pyrenean orogeny. *Marine and Petroleum Geology* 12: 809–820.
- Serrano O, Hanot F. 2005. Le bassin Oligo-Miocène de l'Hérault: un exemple de rétrocharriage des structures pyrénéennes. Implications hydrogéologiques. BRGM. Rapport n° BRGM/RP-53733-FR, 22 p.
- Sigé B, Legendre S. 1997. Un outil de la stratigraphie du Tertiaire continental : l'échelle de niveaux-repères de mammifères. In: Aguilar JP, Legendre S, Michaux J, eds. *Actes du congrès Biochron'97*. Volume Mem.Trav. EPHE Inst. Montpellier 21, pp. 47–54.
- Simons DB, Richardson EV, Nordin CF. 1965. Sedimentary structures generated by flow in alluvial channels. In: Middleton GV, ed. *Primary sedimentary structures and their hydrodynamic interpretation*. Volume Special Publication 12. Society of Economic Paleontologists and Mineralogists, pp. 34–52.
- Sissingh W. 1998. Comparative Tertiary stratigraphy of the Rhine Graben, Bresse Graben and Molasse Basin: correlation of Alpine foreland events. *Tectonophysics* 300: 249–284.
- Southard JB. 1991. Experimental determination of bed-form stability. *Annual Review of Earth and Planetary Sciences* 19: 423–455.
- Teixell A, Labaume P, Lagabrielle Y. 2016. The crustal evolution of the west-central Pyrénées revisited: Inferences from a new kinematic scenario. *Comptes Rendus Géosciences* 348(3-4): 257–267.
- Teixell A, Labaume P, Ayarza P, Espurt N, de Saint Blanquat M, Lagabrielle Y. 2018. Crustal structure and evolution of the Pyrenean-Cantabrian belt: a review and new interpretations from recent concepts and data. *Tectonophysics* 724-725: 146–170.
- Ternois S, Odlum M, Ford M, Pik R, Stockli D, Tibari B, *et al.* 2019. Thermochronological Evidence of Early Orogenesis, Eastern Pyrénées, France. *Tectonics* 38(4): 1308–1336.
- Thaler L. 1962. Premiers résultats d'une recherche systématique des dents de Ronceurs, par lavage de marnes de l'Oligocène, en Bas-Languedoc. *C. R. Somm. Soc. Géol. France* 10: 315–316.
- Tugend J, Manatschal G, Kusznir NJ, Masini E, Mohn G, Thion I. 2014. Formation and deformation of hyperextended rift systems: insights from rift domain mapping in the Bay of Biscay-Pyrénées. *Tectonics* 33: 1239–1276.
- Vandenberghe N, Hilgen FJ, Speijer RP. 2012. The Paleogene Period. In: Gradstein FM, Ogg JG, Schmitz MD, Ogg G, eds. *The Geologic Time Scale 2012*, Volume 2. Elsevier, pp. 855–821.

- Vergés J, Garcia-Senz J. 2001. Mesozoic evolution and Cainozoic inversion of the Pyrenean Rift. In: Ziegler PA, Cavazza W, Robertson AHF, Crasquin-Sleau S, eds. *Peri-Tethys rift/wrench basins and passive margins*. Volume Mémoire 186. Paris : Muséum National d'Histoire Naturelle, pp. 187–212.
- Viallard P. 1987. Un modèle de charriage épiglyptique: la nappe des Corbières orientales (Aude, France). *Bulletin de la Société géologique de France* III(3): 551–559.
- Ziegler PA, Dèzes P. 2005. Evolution of the lithosphere in the area of the Rhine Rift System. *International Journal of Earth Sciences* 94: 594–614.

**Cite this article as:** Séranne M, Couëffé R, Husson E, Baral C, Villard J. 2021. The transition from Pyrenean shortening to Gulf of Lion rifting in Languedoc (South France) – A tectonic-sedimentation analysis, *BSGF - Earth Sciences Bulletin* 192: 27.